Global Warming and its Impact on the Climate in 2024

Editors

Zahraa Radi Harijah Mohammed
Department of Environmental Health, College of Applied Medical Sciences, University of Karbala, Iraq

Aban Talal Mohammed Saeed
Department of Environmental Science, College of Environmental Science and Technology, University of Mosul, Iraq

Saja Abbas Burhi Hussein
Department of Environment, College of Environmental Sciences, Al-Qasim Green University, Iraq

Noor Rasim Yousif Jasim
Department of Environment, College of Science, University of Al-Qadisiyah, Iraq

Hussein Abdul Hassan Talab Haleeb
Department of Environmental Theology, College of Environmental Science, University of Mosul, Iraq

Bright Sky Publications ™
New Delhi

Published By: *Bright Sky Publications*

Bright Sky Publication
Office No. 3, 1st Floor,
Pocket - H34, SEC-3,
Rohini, Delhi, 110085, India

Editors: *Zahraa Radi Harijah Mohammed, Aban Talal Mohammed Saeed, Saja Abbas Burhi Hussein, Noor Rasim Yousif Jasim and Hussein Abdul Hassan Talab Haleeb*

Edition: *1st*

Publication Year: *2024*

Pages: *88*

E-Book ISBN: *978-93-6233-362-9*

Paperback ISBN: *978-93-6233-222-6*

Book DOI: *https://doi.org/10.62906/bs.book.159*

Price: *₹ 537/-*

Contents

Chapter - 1

Introduction

As the earth warms due to the effects of climate change, the resulting shifts in weather patterns have significant ramifications on a global scale. While certain areas may experience a significant decrease in temperatures, the majority of regions will witness an alarming increase in dryness, warmth, or precipitation. The profound and wide-ranging impact of global warming extends to the realm of climate change, giving rise to both perilous and uncertain situations as well as promising prospects across the planet. Amidst these rapidly changing weather patterns, numerous new opportunities arise within the agricultural sector, which has the potential to adapt and innovate to meet the challenges of a shifting climate. However, the potential loss of tropical ecosystems looms as a vast and formidable danger, threatening the delicate balance of biodiversity and ecosystem services they provide.

Moreover, the increasingly rapid thawing of Arctic ice further exacerbates the already rising sea levels, posing a direct and imminent threat to coastal cities and vulnerable communities worldwide. The melting ice not only affects the delicate balance of global climate patterns but also causes significant shifts in ocean currents, leading to disruptive and unpredictable consequences. The loss of Arctic ice also means the loss of critical habitat for unique and specialized species, as well as traditional livelihoods for indigenous communities who rely on the ice for hunting and transportation. Protecting and preserving these fragile ecosystems becomes increasingly crucial in the face of climate change and its cascading effects.

Furthermore, the imperiled state of biodiversity becomes increasingly apparent as various species struggle to adapt or relocate in the face of these rapidly evolving conditions. The pace of climate change outstrips the natural capacity of many species to adjust, resulting in dramatic declines and endangerment. This loss of biodiversity not only undermines the intrinsic value of species but also threatens vital ecosystem functions, such as pollination, natural pest control, and nutrient cycling. It is imperative that we take immediate and concerted action to mitigate the impacts of climate change and protect the irreplaceable biodiversity that sustains our planet.

In conclusion, the consequences of climate change are far-reaching and multifaceted, encompassing shifts in temperature, precipitation, the loss of critical ecosystems, rising sea levels, and the perilous state of biodiversity. The urgent need to address these challenges requires international cooperation, innovative solutions, and a collective commitment to sustainability. By taking bold and decisive action now, we can strive to mitigate the worst impacts of climate change, safeguard vulnerable communities and ecosystems, and create a more resilient and sustainable future for all [1, 2].

Global warming refers to climate change caused primarily by anthropogenic emissions of greenhouse gases. The temperature of the earth has risen by almost one degree since the late 19th century, according to scientific research and evidence. This recent warming is not merely a result of natural internal variability in the earth's climate system; rather, it can be largely attributed to human activities. In fact, there is a staggering 95 percent probability that more than half of the warming observed since the mid-20th century has resulted directly from the actions of humanity.

One of the key contributors to global warming is the burning of fossil fuels such as coal, oil, and natural gas. These fuels, which have long been relied upon for energy production, release substantial amounts of carbon dioxide, methane, and nitrous oxide into the atmosphere when combusted. These greenhouse gases act like a blanket, trapping heat from the sun and causing the planet's temperature to rise. It is important to note that once emitted, these gases can persist in the atmosphere for extended periods, ranging from decades to even centuries, exacerbating the warming effect. Global warming is the increase in the average temperature of the Earth's atmosphere due to the accumulation of greenhouse gases such as carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), and water vapor. Carbon dioxide gas is the most responsible for global warming. It is produced primarily by burning fossil fuels and in smaller amounts by deforestation, certain industrial processes, and agriculture. The second most important global warming gas is methane. It is produced during the use of fossil fuels, the digestion of food by cattle and other livestock, and the decomposition of organic waste in landfills. Other global warming gases include water vapor, nitrous oxide, ozone, and several artificial products known as chlorofluorocarbons (CFCs). The accumulation of these gases in the atmosphere causes the greenhouse effect: part of the Sun’s light is reflected into space, but part of it is absorbed on Earth’s surface, warming it. The Earth cools down at night (and also after sunset), emitting infrared radiation: part of these infrared rays goes back to space, but another part is reflected by the greenhouse gases, increasing the temperature of the atmosphere.

Since 1850, when the first data were collected, the Earth's temperature has risen by about 1 °C (1.8 °F). Most of this warming has occurred since 1975; the increase in temperature has been especially strong in the Northern Hemisphere winter and has strongly affected the Earth's polar regions: the Arctic has warmed about 2–3 °C (3.6–5.4 °F), while parts of Antarctica have cooled. The last decade was the warmest during the last millennium on a global scale, although some regions of the tropics and Siberia have cooled slightly. In the last 30 years, the temperature has also oscillated: after a warm period (1920–1940), there was a cold period (1940–1975) known as "the great global cooling alarm" which spurred controversial theories linking it to the climate cycles driven by the activity of the Sun. However, satellite data and surface observations indicate a monotonic surface warming trend over the last 30 years. The past decade was the warmest on record and all top 10 warmest years occurred since 1990, with 2001 being the warmest year. Global warming has far-reaching consequences for the ocean, the atmosphere, and the land. It will alter the quantity and distribution of freshwater resources and will increase the risk of floods, droughts, and malnutrition.

Global warming refers to the incremental rise in Earth's average temperature due primarily to the greenhouse effect, precipitated by industrial processes, deforestation, and the combustive use of fossil fuels. This triggers changes in the climate system, including alterations to the state characteristics of the oceans, atmosphere, and ice cover, and the distribution of the continents and oceans. It can also affect changes to the Earth's orbits and axial tilt, consequently having either short or long-term effects on climatic conditions.

The greenhouse effect is exacerbated by an increase in concentration of greenhouse gases, including carbon dioxide (CO_2), nitrous oxide (N_2O), ozone (O_3), and methane (CH_4). When solar radiation hits the Earth's surface, a portion is transformed into infrared radiation. This radiation is absorbed by greenhouse gases and re-emitted in all directions. Some of this energy, as a result, goes back towards the Earth, heating the surrounding air. Greenhouse gases thus accumulate at the top of the troposphere, resulting in a heating of its surface and atmosphere. If the concentration of greenhouse gases in the atmosphere increases, then so does the temperature of the Earth as more energy is added to the climate system.

Human activity has caused an increase of greenhouse gases in the atmosphere over the last century. Industrial agriculture, for instance, has transformed surface and buffer lands, logarithmically increasing nitrous oxide emissions from fertilizer use. Deforestation, both with the cutting down of trees and the burning of plantations, increases carbon dioxide emissions. Increased waste production, poorly managed landfills, and gas leaks in oil and gas extraction aggregate methane emissions. Meanwhile, increases in the use of petroleum, coal, and natural gas as energy resources add additional greenhouse gases to the atmosphere. Altogether, human activity has increased total CO_2 concentration by more than 100 ppm, from about 280 to more than 400 ppm, since the dawn of the industrial revolution. This increase occurs alongside a higher concentration of methane, nitrous oxide, and Freon-11. Combined, all the polluting activities have increased the total direct radiative forcing by greenhouse gas emissions by approximately 2.4 W/m2, with a predominantly anthropogenic origin.

In addition to fossil fuel combustion, deforestation and certain land-intensive agricultural practices also play a significant role in contributing to global warming. When forests are cleared, the carbon stored in trees and vegetation is released into the atmosphere as carbon dioxide. Furthermore, the removal of trees diminishes the planet's natural capacity to absorb this greenhouse gas, intensifying its concentration. Similarly, certain agricultural activities, such as excessive use of synthetic fertilizers, can lead to the release of nitrous oxide, a potent greenhouse gas.

As global temperatures continue to rise, the consequences of global warming become increasingly evident. These include but are not limited to more frequent and severe heatwaves, shrinking ice caps and glaciers, rising sea levels, unpredictable weather patterns, and increased intensity of natural disasters like hurricanes and wildfires. The impacts extend beyond the environment, threatening various aspects of human society, including agriculture, water resources, public health, and biodiversity.

In order to mitigate and adapt to the challenges posed by global warming, concerted international efforts are crucial. These efforts involve transitioning to low-carbon and sustainable energy sources, promoting energy efficiency, conserving forests and other natural ecosystems, implementing climate-resilient agricultural practices, and adopting policies that prioritize climate action. The goal is to limit global temperature rise to well below 2 degrees Celsius compared to pre-industrial levels, as agreed upon in the Paris Agreement, and to pursue efforts to limit the increase to 1.5 degrees Celsius. The United Nations Framework Convention on Climate Change (UNFCCC) recognizes anthropogenic (human-induced) climate change as a present-day concern. The accumulation of greenhouse gases in the atmosphere, mainly carbon dioxide (CO_2) emissions from burning fossil fuels, alters the composition of the atmosphere. Global warming is expected to have serious negative effects on ecosystems, organisms, and human societies. Concern for climate change had been talked about in the scientific community for over a century when the industrial revolution began emitting large amounts of greenhouse gases into the atmosphere. The gradual warming of the planet over the following decades aroused further interest and concern. A brief history of global warming leads to where we are today is presented. The background to the scientific investigations of climate change is first discussed, followed by the major international treaty (the UNFCCC) and political discussions concerning the issue.

Until the 19th century, the natural sciences regarded the Earth's climate as a constant. In the early 1800s, Joseph Fourier, a French mathematician, discovered that the average temperature of the Earth was higher than that predicted from its distance from the Sun and the thermodynamics of black bodies, proposing a "greenhouse effect." In 1856, Eunice Foote prepared one of the first scientific experiments to test the effect of increased CO_2 on temperature, making the link between industrialization (fossil fuel combustion) and climate change. In 1896, the Nobel Prize-winning Swedish physicist Svante Arrhenius presented predictions to the Royal Swedish Academy of Sciences. Further understanding of the Earth's climate system and the extent of the human climate influence was obtained in the mid-1900s through the development of meteorological instruments, the first general circulation model (GCM), and what is now called the "overall view of CO_2-forced climate change."

By the late 1970s, climate change led to an increasing interest outside of the scientific community. By the late 1980s, global warming began to enter the public and political domain. Powerful storms, floods, droughts, heat waves, and other abnormal weather events around the world led to speculation in the mass media regarding possible climate change. Along with investigation reports published by worried scientists, these abnormal weather events helped build momentum for the political debate on climate change. In June 1988, then-Environment Minister Kumiko Hasegawa gave the unprecedented policy speech in the Diet, whose first sentence read, "Global climate change is a long-term, grave, and global problem seriously affecting humanity's existence." At the same time, the Intergovernmental Panel on Climate Change (IPCC) was launched by the United Nations (UN) as an intergovernmental scientific panel. The first IPCC Assessment Report (FAR) was published in 1990. In December 1990, G7 leaders agreed at a summit meeting in Paris to the establishment of the UNFCCC. The UNFCCC was adopted and opened for signature at the United Nations Conference on Environment and Development (Earth Summit) in June 1992 in Rio de Janeiro, Brazil.

By acknowledging and addressing the causes and impacts of global warming, and by collectively working towards sustainable solutions, humanity has the potential to shape a more resilient and environmentally conscious future. The urgency to act cannot be underestimated, as the consequences of inaction could be irreversible and have far-reaching implications for current and future generations. It is essential that we embrace the responsibility to safeguard our planet for the sake of not only our own well-being but also that of all living beings who call Earth home [3, 4, 5, 6].

The greenhouse effect is a natural phenomenon that keeps the earth's average temperature at around 60°F (15°C). Without it, the earth would be too cold to support life. Unfortunately, human activity has intensified the greenhouse effect, preventing more heat from escaping, resulting in climate change. This change in climate brings forth a myriad of hazards that pose significant threats to our planet and existence. One such hazard is the increased occurrence of coastal flooding, which poses a tremendous risk to coastal communities, infrastructures, and ecosystems. The rise in sea levels, caused by the melting of polar ice caps and thermal expansion of water, exacerbates this threat further, potentially displacing millions of people and causing extensive damage. Moreover, climate change is contributing to the increased intensity of floods, leading to devastating consequences for communities around the world. The combination of extreme weather events, such as hurricanes, with rising sea temperatures creates a perfect storm of destruction, resulting in widespread devastation and loss of life. Not only does climate change impact coastal areas, but it also has far-reaching consequences for marine ecosystems. Alterations in the salinity of ocean currents, triggered by climate change, can disrupt marine ecosystems, leading to ecological imbalances and the collapse of vital marine industries. The loss of forests, tundra, prairies, and desert shifts, caused by changing weather patterns and habitat destruction, also significantly impact global biodiversity and ecological stability. These changes in ecosystems have a ripple effect, affecting various species and disrupting the delicate balance of nature. It is essential to recognize that while climate change poses numerous challenges, it also presents opportunities for adaptation and innovation. One such opportunity is the potential for increased agricultural productivity in some temperate regions. As temperatures become milder, crop yields can rise, offering a chance to combat food insecurity and improve global food production. This increase in productivity can help feed a growing population and alleviate hunger in vulnerable regions. Furthermore, the melting of polar ice caps opens up possibilities for harvesting Arctic oil reserves, which were previously inaccessible. While controversial due to the environmental risks associated with oil extraction, this opportunity can have significant economic implications. The exploitation of Arctic oil reserves can provide a new source of energy and revenue, creating jobs and stimulating economic growth. However, it is crucial to approach these opportunities with caution and ensure that sustainable practices are in place to minimize the negative impacts on the environment. Overall, addressing climate change requires a multi-faceted approach that considers both the threats and opportunities it presents. It is crucial that we focus on implementing sustainable practices, mitigating the impact of climate change, and exploring these opportunities, all while prioritizing the preservation of our planet and its delicate ecosystems. Only through collective action and global cooperation can we tackle the challenges of climate change and pave the way for a brighter, more sustainable future [7, 8, 9].

1.1 Background on global warming

Global warming is the gradual and persistent rise in the average temperature of Earth's atmosphere and oceans. This unprecedented phenomenon is primarily driven by human activities, particularly the burning of fossil fuels and deforestation, which release significant amounts of greenhouse gases into the atmosphere. As these greenhouse gases accumulate, they form a thick layer, trapping heat from the sun and causing the planet's temperature to steadily increase.

In recent decades, global warming has become a pressing global concern, attracting widespread attention from scientists, policymakers, and the general public. The magnitude and urgency of this issue cannot be overstated. The consequences of global warming extend far beyond simply warmer temperatures. They manifest in a wide range of challenges and disruptions that affect Earth's delicate ecosystems, weather patterns, and human societies.

One of the main contributors to global warming is the burning of fossil fuels for energy production, transportation, and industrial processes. As we continue to rely heavily on these finite energy sources, carbon dioxide and other greenhouse gases are released into the atmosphere on an unprecedented scale. Moreover, deforestation exacerbates the issue by reducing the Earth's capacity to absorb these emissions, further amplifying their warming effect.

The consequences of global warming are already being felt worldwide. Rising sea levels, intensified heatwaves, more frequent and severe storms, and prolonged droughts are just a few examples of the far-reaching impacts of climate change. These environmental changes pose significant threats to biodiversity, food security, water resources, and public health, affecting both natural ecosystems and human communities.

However, it is crucial to recognize that the consequences of global warming are not evenly distributed. Marginalized and vulnerable populations, including indigenous communities, low-income countries, and small island nations, are disproportionately affected. These groups often lack the resources and adaptive capacities to cope with the impacts of climate change. As a result, their livelihoods, cultural heritage, and even survival are under severe threat.

Addressing global warming requires international cooperation, decisive actions, and transformative changes in various sectors of society. It is essential to reduce greenhouse gas emissions by transitioning to renewable and sustainable energy sources, improving energy efficiency, and adopting cleaner technologies. Additionally, efforts should focus on enhancing resilience and adaptation measures to minimize the social, economic, and environmental impacts of climate change.

Ultimately, combating global warming is not only an environmental imperative but also a moral obligation to safeguard the planet for future generations. By acknowledging the gravity of the situation and working collectively, we can mitigate the effects of global warming and build a sustainable and resilient future for all.

Climate is defined as the long-term change in temperature, rainfall, humidity, winds, and seasons in a particular region. It is affected by both natural agents and human activities. Global warming, air pollution, and climate change are common scientific concepts discussed across the world. The nation has been affected by climate change and rising sea levels, resulting in challenges for government, land, and agriculture. Scientists and governments constantly monitor the climate to find ways to tackle the impacts and changes in climate.

Global warming is caused primarily by the excessive emissions of greenhouse gases that result from the burning of fossil fuels, the intensive practices of agricultural production, and the alarming rates of deforestation around the world. These detrimental emissions are a direct result of our everyday activities, such as using cars for transportation, relying heavily on electricity for our needs, and employing various modes of transportation. Sadly, the increasing dependence on fossil fuels and the widespread use of forests for agricultural purposes have drastically heightened the carbon output, resulting in a surplus of greenhouse gases within the atmosphere.

Consequently, these greenhouse gases become trapped, disrupting the natural cycles that regulate our planet and ultimately leading to the entrapment of heat. While climatic changes have historically occurred in a natural manner, the current and continuous ascent of greenhouse gas levels is undeniably unnatural and can largely be attributed to the activities of humankind.

Climate change is unequivocally and indisputably one of the most paramount and pressing human development challenges of this century. In every corner of the globe, vulnerable communities and nations bear the brunt of the adverse effects caused by climate change, resulting in an alarming and exponential escalation of water scarcity. This dire situation amplifies the risks and perils faced by millions, leading to a substantial loss of livelihood and a bleak future filled with profound uncertainty.

The consequences of climate change are far-reaching and profound, as it gravely diminishes economic growth prospects for nations near and far. Objectively speaking, the ramifications are dire and cannot be underestimated or ignored. The mere existence of a minuscule increase in temperature can no longer be seen as inconsequential. On the contrary, it has the potential to precipitate catastrophic consequences of unfathomable magnitude.

One of the most disheartening aspects of climate change is its insidious and devastating impact on the delicate ecosystem that sustains our planet. The relentless march of global warming threatens to unravel the intricate web of life on Earth. Countless plant and animal species find themselves teetering on the brink of extinction, as their habitats become compromised and their chances of survival diminish with each passing moment. This catastrophic unraveling has the potential to irrevocably alter the natural balance we have come to rely upon and cherish.

The significance and seriousness of climate change cannot be overstated or denied. Countries that have been ravaged by its merciless effects bear witness to the immense destruction and anguish it brings. Bangladesh, a nation constantly grappling with the devastating impact of rising sea levels and extreme weather events, stands as a somber testament to the urgency and gravity of this paramount issue. Similarly, Japan, a country known for its resilience, has experienced firsthand the ruthless power of climate change through frequent and devastating natural disasters.

The science of global warming

Introduction to global warming

The greenhouse effect is a natural phenomenon that warms the Earth’s surface. When the sun’s energy reaches the Earth, some of that energy is reflected back to space and the rest is absorbed, warming the Earth. The Earth, in turn, radiates energy back to space in the form of infrared, or heat, radiation. Since some of the Earth’s surface is relatively cool (average of around 15 degrees Celsius), this radiation has a long wavelength (about 10 microns). Greenhouse gases are gases in the atmosphere that can absorb this radiation, subsequently emitting some of this energy again back to the Earth’s surface. Greenhouse gases include water vapor, carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O). Greenhouse gases effectively trap heat at the Earth’s surface. This re-absorption of heat has a warming effect, which is essential for the Earth’s climate. However, in the absence of greenhouse gases, the Earth would be around 33 degrees Celsius cooler.

The effect of human activities on greenhouse gases has been the cause of an increasing warming effect, subsequently leading to the enhancement of the greenhouse effect and global warming. While there was a relatively constant concentration of CO_2 due to a balance between natural phenomena in the preindustrial period, man-made activities (e.g. fossil fuel combustion and other anthropogenic activities) have led to an increasing concentration of CO_2. A similar development can be seen for CH_4 and N_2O. Since the mid-19th century, the concentration of CO_2 has steadily increased from about 280 ppm to almost 400 ppm (currently around 380 ppm), while the concentrations of CH_4 and N_2O have greatly increased as well. Greenhouse gases retain heat very efficiently. As a result, even small changes in their concentration in the atmosphere can lead to considerable climate effects. It was estimated that a doubling of the preindustrial CO_2 concentration would lead to an increase in the average global temperature of about 1.5 to 4.5 degrees Celsius.

Sunlight travels through space to reach Earth, where some of it warms the ground and ocean. Bodies of water and the Earth's surface absorb quite a bit of sunlight energy, so they become warm. Sunlight primarily arrives in the form of visible light at the top of the atmosphere, where some is reflected by clouds and the surface, while the vast majority reaches the ground. "But the warm Earth doesn't just sit there passively waiting for the next beam of sunshine," scientists might exclaim. "It radiates energy back to space in the form of infrared radiation (heat). If it didn't, the Earth would continue to heat up until it became so hot that all available water turned to vapor." Everyone agrees this is not in the cards for our home planet, so why is the average temperature rising?

Unlike visible light, infrared radiation is not invisible to greenhouse gases in the atmosphere. These remarkable molecules (mostly H_2O and CO_2) vibrate and turn, absorbing infrared radiation for a short time while then radiating it back in all directions, with at least half of it returning towards the surface. This lovely arrangement is just the atmospheric greenhouse effect that saves many Earth-like planets from being frozen inhospitable ice worlds, keeping the average temperature on Earth a mild and comfortable 15°C (almost 60°F) instead of a frigid −18°C (or 0°F) as predicted by its black body temperature. But... "Wouldn't this mean that more greenhouse gases in the atmosphere would lead to greater warming, which might eventually spiral out of control?" scientists wondered. The balance between energy arriving to earth and energy escaping into space is: luminosity of the sun (Lsun) = 3.84×10^{26} Watts = 341 W/m² times cross-sectional area of the Earth (Af) = π x R² = $\pi(6.37 \times 10^6)^2$ m² or 1.276×10^{14} m²; this leads to a total of 341 W/m² of energy arriving to the Earth which is equal to the net average over the surface of the earth of 239 W/m² energy escaping to space calculated from earth's radiative cooling.

Changes in temperature, precipitation, and sea levels, and other Earth systems are considered indicators of climate change. These indicators reveal the difference between "natural" climate variability and human-induced climate change. They are tracked by scientists all around the world and summed up in a periodic report, but there are several fundamental indicators that keep track of climate without a time cap. The most significant of these are temperature changes and sea level rises.

Temperature changes

Temperature changes are possibly the most well-known climate change indicators. Many people are aware of how the average global temperature has increased by around 1.2°C from the late 1800s to the early 2020s. This increase in temperature has long-standing effects on natural systems and has begun to severely affect human systems. Temperature indicators reliably predict climate change because of a good understanding of natural variability on many timescales by scientists. There are two essential components to climate sensitivity. The first is how much additional warming is predicted as atmospheric CO_2 concentration increases. The other is how easy it is to return the climate to the original state. Global temperature averages are commonly regarded as the most important summary measure of climate, similar to how GDP is commonly used to measure the economic performance of countries.

Sea level rise

Sea level rise is more problematic in climate terms than temperature, but it remains one of the most important climate change indicators. Sea level changes reveal climate changes over the whole earth system, including the atmosphere, oceans, and cryosphere, but not over land systems like temperature. Different timescales of climate variability also influence sea level, making the attribution of a change to human influence troublesome. Predictions for future sea level rise have higher uncertainty than for temperature. The most complex systems still predict temperature changes over several centuries but only freeze the large Antarctic and Greenland ice sheets on a timescale of millennia. Still, sea level rise is the second most scientifically robust climate change indicator after temperature changes. It has grown more certain in the early 21st century, approaching an annual rise of 1 cm in 2007, which exceeds what was expected in IPCC 2001 (0.1-0.9 cm/yr).

The average global temperature has increased by nearly one full degree Fahrenheit since the late 1800s. That may not sound like much, but it has profound effects on the world. The rising temperature is mainly due to increased carbon dioxide in the atmosphere from burning fossil fuels. The Earth's average temperature is easier to understand, since the Earth has a hot side and a cold side. The hot side faces the sun and generally matches the surface temperature of Venus, while the cold side cannot be seen from Venus and generally matches the surface temperature of Mars. Because both Venus and Mars are not blanketed in carbon dioxide like Earth, their average temperature is consistent around both day and night sides.

The Earth's atmosphere is about 300 miles thick. Since the Earth is 8,000 miles in diameter, that means the atmosphere is very thin when viewed from the side, like the skin of an orange. The moon has no atmosphere, so when the sun goes down, temperature on its surface falls to minus 280 degrees Fahrenheit. The average temperature on its dark side is minus 400 degrees Fahrenheit. This is approximately the surface temperature of the Earth if it had no atmosphere. However, with the Earth's atmosphere, its average temperature is not 400 degrees Fahrenheit, but rather a balmy 59 degrees Fahrenheit. This is a very comfortable temperature for us and is maintained when the Earth's icecaps are not melting. By covering a planet with the correct amount of carbon dioxide, the average temperature can be brought to the point that liquid water can exist.

Temperature changes have, can, and will occur naturally over geologic time. Changes in solar output, twist and turns of the Earth's orbit and tilt, meteor impacts, super volcanoes, breaking up of continents forming new ocean basins, and the slowing of the Earth's mantle all change the amount of heat on the Earth. However, the warming from burning fossil fuels, all of which are stored carbon, is up to 50 times faster than any natural warming. One of the reasons temperature change appears gradual is that there is a considerable lag from when heat is added to the climate system and when it is felt on the Earth's surface. The ocean takes a lot longer than the atmosphere to warm, and because burning fossil fuels has been warming Earth for over 250 years, many of the worst impacts of climate change are still yet to come.

To the average observer, it may seem that the sea level has remained constant for many years. In fact, there are many historical references to specific objects or landmarks "by the river" or "in view of the sea." However, over the centuries, many of those objects have ended up underwater as the sea level has risen. This rise of the sea level is still currently measured today and has been identified as one of the most robust indicators of climate change. Similar to the land, the sea level also does not have a perfectly smooth surface.

In the Earth's coordinate system, the sea level can be defined as the mean ocean surface based on the average high and low tides. But as a result of many tides and variations in atmospheric pressure, the local sea level varies daily. Although the sea level may vary locally with respect to this mean ocean surface, the average sea level is about 370 meters higher at the equator than at the poles. This is a consequence of the bi-oidal shape due to the Earth's rotation, the complicated gravimetrical field, the interaction with ocean currents, the atmosphere, and many more.

Before measurements of the absolute sea level began, these tides were measured from a fixed point on land at locations along the European coast where, for instance, the building of a new lighthouse was planned. These measurements were made for several years with the introduction of measuring devices worth several months of work from expert teams. Each measuring point had to be carefully selected away from harbors and rivers and with a solid basis of hard geologic material. Also, during the period of measurements, strict rules applied to care for the devices and their surroundings to avoid "noise" like running a mill or moving large objects across the ground in the vicinity of the measuring devices. The need for good measurements and the extensive preparation of this effort indicates that people were aware of "the sea" and its possible variations that could affect their livelihood, like flood risk when the sea level rose.

The absolute rise of the sea level is attributed to three main causes: thermal expansion, melting of land ice with glaciers and ice-caps, and runoff of snow and ice from glaciers. Thermal expansion responds to increases in temperature of the ocean. As the temperature increases, the average energy of the water molecules will increase, which will eventually result in an increase in volume and consequently a rise of the sea level. Two satellite missions have been launched to derive the absolute sea level: Topex/Poseidon (1992-2007) and Jason 1 and 2 (2001 and 2008 respectively). Since 2006, the satellite mission Jason 1 has been collecting data together with the in-orbit co-operations such as a Ku band altimeter measuring the sea surface height and a radar altimeter system that is used to derive the significant wave height. These on-board systems are supported by a precise GPS-based orbital determination system which allows estimations of location deviations in the order of 1 decimeter as well as a precise trajectory modeling, ground receiving stations with antennas, radar precision timing and caretaking units, and signal preparing laboratories.

Global warming is the increase in the Earth's temperature caused by the buildup of greenhouse gases in the atmosphere. Personal actions, like driving cars and heating homes with fossil fuels, add to the buildup. So does the less visible burning of forests and clearing of land for agriculture. It was first predicted over a century ago; since then, the burning of fossil fuels and deforestation have grown rapidly, and global temperatures have increased. It begins by describing how ecosystems like forests and coral reefs are being severely damaged in response to rising temperatures and their consequences on the existing biodiversity. It then goes on to explain how these changes in the environment are also affecting human health.

Rising temperatures are damaging ecosystems like forests, coral reefs, and tundra. Coral reefs are dying and becoming bleached, as the algae that live within them die off in high temperatures. Coral reefs are not only beautiful, but they also support a rich biodiversity, and once lost, these reefs will be difficult to restore. Warming has caused the survival range of some species, like the cold-water fish, to decrease, potentially resulting in their extinction. In diverse ecosystems that have existed for thousands of years, such as tropical rainforests, temperature-induced stresses will create competition between species. Non-native species that are better suited to higher temperatures will replace or expand into similar habitats at the expense of vulnerable native species, thus reducing biodiversity. Evolving rapidly to adapt is difficult for long-living species like trees; and when the evolution happens, they may no longer fit in an existing ecosystem. Climate change would create a "winner-win" situation for some species that have adapted and a "loser-lose" situation for highly specialized ones similar to "fast and slow."

Distributed across continents, nations, and communities; with an already tiring burden of poverty, hunger, illiteracy, lack of equal development, and politically-correct action; and unaware of unmanageable environmental change, the relatively poorer and vulnerable group stricken by problems is stupendously devastated by global warming. There is a drastic change in weather and rainfall patterns for various places over each continent, adversely affecting agricultural practice. A gradual increase in dry and barren lands will occur in most tropical areas, thus lowering food production, devastating the subsistence of the poorest nations. With a high-density population and low countries, for these nations, havoc floods are caused by rising sea levels and storms, as is seen in Bangladesh, slowing their development efforts. Along with low-lying coastal and island regions, the Western Pacific is especially threatened, imposing a national security dilemma. Harmful migratory and rescue problems will be faced, the poorest always entering the victim's course first. Depending upon personal actions, rising greenhouse gas concentrations are foreseen, producing a rise in medical problems.

Global warming triggers a cascade of destructive changes in ecosystems, irrevocably altering their physical, chemical, and biological components. The Intergovernmental Panel on Climate Change (IPCC) expects a dramatic impact on the environment and biodiversity. Many populations of terrestrial, freshwater, and marine species are shifting their distributions towards higher latitudes and elevation due to climate conditions. Ongoing stressors, such as habitat loss, pollution, and invasive species, further increase organisms' vulnerability to climate change. Globally, reconstructed empirical data indicate that the biomass of plankton organisms has decreased in the ocean's upper valid portions through 40 years of warming. Coral reefs, home to 1 million marine species, are imperiled by warmer waters resulting from climate change; 19% of coral reefs have died in the last 50 years. Harmful algal blooms are increasing due to the combination of warmer water, nutrient runoff from land, and complex interactions that generate increased emphasis on harmful species. The growing threat of harmful estuarine and coastal algal blooms is anticipated. Recent research has shown that potential climatic changes, focusing on rainfall, temperature, and oxidation state, can greatly affect concentrations of phosphorus (P) in sediments and dissolved oxygen (DO).

The negative effects of climate change on terrestrial ecosystems are critical and complex, involving influences on components and processes such as forest growth, biomass accumulation, soil processes, hydrology, and climate itself. Studies with very different methodologies, data quality, and representativeness are analyzed and summarized to glean general patterns and estimates of the sensitivity of terrestrial ecosystems to climate warming. Changes in climate and atmospheric carbon dioxide (CO_2) concentration influence ecosystems' structure and functioning through alteration in vegetative growth, primary productivity, decomposition, respiration, soil processes, and loss and gain of carbon, nitrogen, and moisture stores.

Inaccurate equations in some models can continuously alter soil moisture, anomaly temperature, and productivity over areas, which easily induce significant variability for carbon exchange among unreasonably large areas. A heat uptake and disequilibrium between energy contents of ocean-lithosphere-hydrosphere-atmosphere are provided by the input CRM coupling to the German BMBF 10-year project AWAKE.

Soil N availability and moisture are induced by the coupling between a heat source anomaly and disequilibrium expansion of the ocean. Changes in the seasonal and geographical distribution of climate loading and land *et al*. patterns of N flux and moisture are induced by changes in the mean ocean haline stratification. Through microscopic modelling of the physical interface, a system of pressure equilibrium is developed to model the "starving lens" generator, broadly applicable to frictional zones. Systematic reference solutions and some performance measures are provided to guide the calibration and use of the models developed for climatic studies. New shared data are created that allow modelling groups to efficiently assess spatially explicit, long-term effects of climate change on N cycling.

The most serious threat posed by global warming may be to human health. Indeed, rising temperatures and shifting climatic patterns are likely to have numerous harmful effects on people, including increased rates of heat-related deaths, exacerbated air pollution, increases in some infectious diseases, and changes in the distribution and life cycles of vectors that transmit these diseases. In addition, warming may produce food and water accessibility problems.

In 1986, a heatwave and subsequent drought hit large parts of North America and the Soviet Union. In the United States, average temperatures were 2 degrees Celsius above the historical norm, and deaths from heat exposure rose sharply in many cities. In Chicago, the number of deaths attributed to the heat was about seven times higher than in previous "normal" years. The elderly and chronically ill were particularly vulnerable. Globally, weather-related mortality due to heatwaves was sharply up, from about 100 people killed in the average year in the 1950s and 1960s to over 2,115 deaths in 1991 and more than 66,603 deaths in 1995.

Many other cities in the Northern Hemisphere also experienced similar events in the 1980s and 1990s. The index of deviation of monthly temperature from the long-term mean (ANOM monthly temperature) for the months July, August, and September of 1988 indicated a linear trend of 1.57°C per year. Yet another North American heatwave struck in 1995, from July 11-29, pushing heat index levels above 115°F around central U.S. states. The mean temperature of the contiguous United States from the January-July period in 1995 was estimated at +1.68°C relative to the 1961-1990 base, and cities on the Great Lakes were most severely affected.

Human health effects of environmental change are mediated through changes in biogeochemical, hydrological, and ecological processes. Changes to environmental changes, temperature, precipitation, water quality, and water quantity are found to directly affect human health.

Contemporary policy interventions must ensure the use of renewable energy sources at all stages of electricity production, starting from the global level (resolution of the UN General Assembly concerning prevention of the catastrophes of climate change can be considered the first step) and finishing with the level of individual countries (energetic policies of the European Union can be considered quite effective examples).

On the scale of as big as possible power stations, the use of water, wind, and waves for electricity production must be planned. The solution to these problems can be the joint projects of several countries (for example, on the mutual border or on the big rivers as in the case of North Hydropower Cascade projects of Russia and Kazakhstan connected with the construction of Zhanadarya and Baype power stations; the projects of joining water channels of the Caspian Sea and Dead Sea are under consideration). The attempt of technological packages exportation should be done. Such experience does exist (as Oil-for-food program flowing from the embargo of western democratic countries against Iraq). At the middle level of power stations, technologies to produce prosperous bio-fuel from wood waste, straw, wax, and other wastes must be used. The use of such technology by Russia, Scandinavia, and Canada can be considered. Small power stations using the regenerative sources must be supported at the national levels (for example, Russia can take use of using atomic batteries powered by the waste of atomic reactors).

At the level of apartments and one-house estates, the policies of using solar batteries, heat counters, fuel cells, and wind power generators must be performed (The Carter solar project of the USA, incentive systems of using wind power in Denmark, domestic stove in Holland, batteries supplying for individual houses in Germany). To avoid the storm of environmental disaster, the mutual actions of countries will help with the question of future generations' existence. It is impossible to reduce the emission of greenhouse gas in equal proportions everywhere. The first steps in the sphere of hard law must be done. There is a prospective sanction of various types for the countries over the norm of permissible emissions, as for the countries below the norm (dropping off developing countries' development rights on the provision of compensations). Environmental issues are getting more and more political significance. At world summits, environmental problems passed from the sphere of platform discussions (as Dublin summit in 1994) to the sphere of frames of concrete programs realization (Kyoto protocol was ratified by developed countries, something like a statement of intentions, presentation of best wishes, and requests to behave reasonably). More importantly, the countries outside of "noble fate contagion risk (to be really more serious and possible enough to perform at the present moment)" must be controlled either. For that, cooperation among the countries of different "noble fate" must be powerful enough (being remembered that global thermal catastrophe only affects developed countries and states). A close cooperation between scientists and practitioners/climatologists/biologists/economists in the sphere.

The rapid depletion of natural resources and energy crises have propelled a global quest for renewable energy sources. Scientific and industrial advancements are fostering the development of efficient and eco-friendly energy production processes. Renewable energy sources can potentially restore the natural balance, resulting in non-polluted energy generation. They involve relatively low operating costs and are free from waste disposal issues, nuclear hazards, and regional energy use conflicts. Popular renewable sources include solar, wind, biomass, hydropower, and geothermal energy.

Solar energy, the most abundant renewable energy source, is harnessed through solar collectors that capture sunlight and convert it into thermal energy. Passive solar energy systems store energy without mechanical intervention. Wind energy is another well-established and widespread renewable energy source that utilizes wind turbines to convert airflow into electricity. Wind energy is free, clean, and inexhaustible, with horizontal-axis wind turbines (HAWT) and vertical-axis wind turbines (VAWT) from which the former is more efficient.

Biomass refers to the organic matter produced by plants and animals, including solid, liquid, gas, and a combination of these states. The biomass utilization processes include combustion, gasification, pyrolysis, and fermentation. The biomass provides energy in the form of electricity (cogeneration), steam, hot water, synthesis gas, bio-oil, and biofuels.

Hydropower refers to the gigantic potential energy of water, being a traditional source of energy, is renewable, and is of low cost compared to other forms of energy. Large hydropower stations can produce electricity in gigawatts (GW). However, these installations can disturb the balance in the ecosystem by submerging large tracts of forestland.

Geothermal energy is harnessed from the internal heat of the Earth. The energy is produced from hot steam trapped in volcanic rocks, fissures or underground aquifers in the earth's crust developed due to the natural decay of radioactive isotopes in the rocks. Geothermal power plants have a minimum carbon footprint. However, these plants can lead to increased acidity, land subsidence, and a lack of dioxides in the crust can result in the premature seals of reservoirs.

Amidst the rising challenge of climate change and global warming, nations and local regions are coming up with creative ideas that might help. Some of these ideas are being considered or tried out, while others have already been established in several places. Overall, the focus is on policy interventions that change the world in a positive direction. However, some consider these changes unfriendly, unnecessary, economically expensive, and impractical.

The carbon pricing policy aims to make polluters pay for their emissions. Under this policy, those who want to emit carbon will need to hold permits demonstrating that they have the rights to emit a certain amount of CO_2. This record of rights can be bought and sold, meaning that emissions allowed are not set in stone. The core principle is that polluters will incur costs that depend on how much they choose to pollute. Therefore, those who target sustainability are preferred to those who continue polluting unnecessarily.

Carbon pricing policies have already been successful in some countries or regions. British Columbia is a province in Canada implementing this policy from July 2008. Under the policy, a tax on fossil fuels increasing every year aims to limit carbon dioxide emissions. Rio de Janeiro is currently running trials of a similar policy to foster sustainable practices. The lagoon and sewage's untreated nitrogen dumping program incentivizes producers to purify their waste before dumping.

Going a bit further, the Clean Air Act Amendments established a cap and trade on sulfur dioxide in the US, solving acid rain pollution at a much lower cost than other alternatives. A similar experiment was done with chemicals thinning the ozone layer, where some of them were eliminated. The 1987 Montreal Protocol aimed to stop creating new ozone-depleting substances and the production of existing ones, bringing new communities resources and spurring research and industrial investment.

Investing in scientific knowledge dissemination, technology, infrastructure, and social organization focusing on sustainability is better than punishment policies. For example, the pay-as-you-throw collection, where fees are based on the number of bags thrown away by residents, promoted recycling and reduced waste levels in several municipalities. Other communities offered rewards to encourage recycling.

Global climate models (GCMs), also known as global circulation models or simply climate models, are numerical representations of the interactions among the atmosphere, oceans, land surface, and ice. Earth's climate system is incredibly complex, with many components dynamically interacting over a vast range of time and space scales. Models can assist in understanding this interconnectedness by simulating what would happen to some characteristics of the system if changes occurred in others. Stratified in research areas, GCMs play increasingly extensive roles in climate-related work scopes, such as long-range forecasting and climate change analysis.

Over the years, advancements in technology, satellite, and computing have resulted in increases in data availability and model resolutions. Climate models have, thus, evolved significantly, facilitating the assimilation of diverse data sets and further exploration of indispensable climate variables. Widely characterized as three-dimensional, deterministic, time-dependent, nonlinear computer models, GCMs solve the governing differential equations by averaging and transcribing them into macroscopic equations. Data normalization via parameters may be controversial, with inadequate resolution on spatial variability. As such, modeling by assembling the output of smaller climate models simulating components of the system also qualifies as GCMs.

Modeling applicable spatial and temporal scales of interest can either prove subtle or stringent, but early studies still provided forecasts ten times the length of the model climate. Moreover, it is dubious whether experimentalists would run up to 100 different models on global decadal time scales. Still, GCMs are powerful climate simulation tools that are better than nothing. Relying on GCM output in support of decades-long planning is not unreasonable, yet the interpretation of risk needs to be stringent as even a few degrees could cause a food crisis half as bad as one following the end of the Little Ice Age.

The basic components of a world model, primarily numerical calculations, the required meteorological quantitative data, the modeling of subsystems, and useless or unnecessary modeling, are presented in this section. The effects of scale, boundary conditions, and mathematical description are discussed. The basic components of a world model are numerical calculations, on the basic processes in the climate system, consisting of the different subsystems (ocean, atmosphere, land, sea ice, etc.); on interactions between subsystems; on the combination of processes and interactions in a global system; on the global energy balance, and on the changes in the system with respect to time.

To calculate the above, required meteorological quantitative data are divided into groups of subsystems and the associated models. A pool of the best modeling, valuated issues (at least, output includes anticipated climatic change), and a model of the natural climate can be needed. Basic processes are surface water balance, energy processes (both radiative balance and heat transfer processes), energy storage, changes in ice areas, land subsystems, mix layers, restoration, circulation, near-surface processes, internal processes on waves, turbulence and diffusion, and biological processes. The modeling of subsystems is on the ocean and its regions; on the atmosphere, regional details on feedback mechanisms (vegetation, sea, and land ice); on land; on spatial scale; and on the simulation of natural variability in a simplified global climate model to investigate climate variability on decadal time scales.

Modeling of useless (or as much as possible simplified, but necessary) or unnecessary modeling includes atmosphere and ocean thermal tides; daily without ridging snow; daily without pressure; daily without clouds; annual average-albedos; and ocean without gyres (only large-scale thermohaline circulation). There are four aspects of the modeling of the ocean to put (as much as possible) in the presented climate model. These aspects are the global bathymetry (contour of ocean bottom) of a model grid, daily temperature, and salinity (as much as possible) to relate the barotropic model sea level with the surface density, temperature, and salinity from the climatological world model; and a more simple parametrization of the wind stress on ocean topography and representation of western boundary currents.

Climate change has been a contentious issue for decades, provoking heated debates. Despite scientific consensus regarding its anthropogenic origins, skepticism has remained a thorn in the side of climate action. Why are some people skeptical? Several factors are at play: political affiliation, social identity, and misinformation can reinforce or build skepticism. Here, the climate skepticism controversy is examined. Leading skeptic arguments are discussed in the context of the scientific consensus, and effective scientific responses to combat climate skepticism are outlined.

Misconceptions about climate change have persisted despite nearly universal scientific consensus. Sometimes classified as myths, false statements about climate change or climate science warrant further exploration. The credibility of those who promote the "myths," questioning the sincerity and motives of climate skeptics, has garnered renewed attention, especially amidst attempts to convey scientific consensus. However, myths cannot simply be dismissed via fact-checking, as they can undermine understanding of scientific consensus. Addressing common concerns or misconceptions surrounding the ongoing scientific consensus can, nonetheless, lead to more constructive dialogues about climate change.

In conclusion, the challenges posed by climate change are daunting and demand our immediate attention and action. The future of our planet and the well-being of current and future generations hang in the balance. It is essential that every nation, community, and individual recognize the urgent need for comprehensive and sustainable solutions. By coming together and prioritizing environmental preservation, we can strive to mitigate the effects of climate change and safeguard the precious resources that sustain us all. Only then can we secure a brighter and more resilient future for ourselves and the myriad of species that share this beautiful planet with us [10, 11, 12].

Chapter - 2

The Science of Global Warming

The Earth's climate is, for every living organism, an essential system of vital importance. Indeed, it would not be possible to understand the development of humanity in the absence of variations in the atmosphere, hydrosphere, cryosphere, and biosphere of the Earth if we analyze the periods that preceded the Holocene. Unfortunately, people often think that global warming is an invention of scientists or that its consequences are so removed in time that the fight against this phenomenon is prevented. Both the former and the latter are false. The climate forms part of the environment of the human being, intimately conditioned and modified by him. What is being carried out is a large-scale weather modification.

Early in this century, these warning voices began to take on an increasingly significant and influential internationally professional dimension. The profound and stimulating literature, which is filled with deep emotions and captivating ideas, now occupies an exceedingly vast amount of space in the most diverse sections of libraries and book collections worldwide. The treatment and understanding of the global warming phenomenon from the scientific point of view comprises an extensive body of knowledge derived from numerous branches of science, including but not limited to atmospheric physics, paleoclimatology, meteorology, oceanography, biology, ecology, geology, astronomy, and countless other interconnected disciplines. It is indisputable that a complex and multifaceted theme like global warming necessitates the incorporation and integration of the invaluable insights and findings from these diverse fields of study. This integration allows for a comprehensive analysis and interpretation of the intricate mechanisms and intricate processes that contribute to the daunting challenge of global warming mitigation. As research continues to advance and our understanding deepens, the urgent need for collaboration among scientists, policymakers, and society as a whole becomes increasingly evident. Together, we must strive to develop innovative solutions and implement effective strategies in order to safeguard the planet's future and mitigate the adverse impacts of global warming on our environment, ecosystems, and ultimately, humanity itself [13, 14, 15].

2.1 Greenhouse gas emissions

As a direct consequence of the remarkable strides made in economic development and the notable transformations in consumption and production patterns, the incessant surge in electricity demand persists with unwavering determination. From an economic standpoint, it is an indispensable element in contributing to bolstering economic growth and the overall welfare of numerous nations, serving as the lifeblood of progress and prosperity. Simultaneously, the utilization of electricity for power generation has exacerbated the escalation of carbon emissions, releasing echoes of environmental concerns in every corner of the globe. Consequently, the urgent need to address air quality concerns and curtail greenhouse gas (GHG) emissions has become a paramount, non-negotiable priority for every one of the 100 cities across the vast expanse of our planet.

Hence, international treaties such as the Kyoto Protocol have diligently enforced regulations pertaining to GHG emissions, heralding a new era of environmental conscientiousness and accountability. The colossal magnitude of GHG emissions is, at present, regarded as a pivotal aspect in formulating a country's economic growth plan, intertwining the intricate fabric of sustainable development and global stability. Governments, corporations, and non-profit organizations alike have united in fervent solidarity, mobilizing their vast array of resources and embarking on an arduous yet essential journey to effectuate substantial reductions in GHG emissions. With resolute determination, they actively embrace the expanding repertoire of green energy solutions, as the future of our planet and subsequent generations lie in the balance.

The green energy sector, propelled by commendable efforts and fortified by governmental subsidies and the unwavering operation of market mechanisms, has experienced a momentous upsurge in unparalleled prominence and influence, surging forth as a beacon of hope amidst the uncertain tides of change. It stands as a testament to human ingenuity and unwavering resilience, providing a tangible pathway towards a greener and more sustainable future. The winds of change have swept across the land, as solar panels stretch towards the sun, wind turbines dance in harmony with the relentless gusts, and hydroelectric power harnesses the raw power of mighty rivers.

Together, we embark on a transformative expedition, our collective footsteps guided by the unwavering compass of environmental consciousness. We embrace the invaluable lessons of the past, drawing inspiration from the mistakes and triumphs of yesteryears, weaving them into the tapestry of progress and innovation. Let us foster an unwavering commitment to sustainability, for within its embrace lies the key to a world brimming with untold possibilities. Let our actions reverberate through time, creating a legacy of stewardship and responsibility for generations yet to come. The greener tomorrow we long for is within our grasp, awaiting our unwavering resolve and collective efforts to usher it into fruition. Together, we march towards a future imbued with the radiance of renewable energy and the boundless potential of a planet in harmonious coexistence with its inhabitants [16, 17, 18].

Greenhouse gases such as carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O) have the significant effect of reducing the acceleration of Earth's radiation, ultimately leading to a notable increase in Earth's overall temperature. It is essential to comprehend that greenhouse gases possess unique characteristics wherein they absorb Earth's radiation at a specific wavelength to a high extent, while simultaneously emitting and reflecting almost all incoming solar radiation wavelengths.

Human economic activities play a pivotal role in the emission of these primary greenhouse gases. CO_2, for instance, is predominantly generated from burning fossil fuels to fulfill various energy needs like electricity generation, heating, and transportation. On the other hand, CH_4 emissions are mainly derived from sources such as rice fields, agricultural and industrial activities, landfills, and waste treatment processes. Similarly, N_2O emissions primarily originate from the application of nitrogen fertilizers, fossil fuel combustion, and deforestation.

It is worth noting that greenhouse gases persist within the atmosphere for a considerable duration, ranging from 100 to 100 years. Notably, these gases possess a higher global warming potential than CO_2, making their impact on the environment more pronounced and concerning. Understanding the intricate nature of greenhouse gas emissions and their prolonged atmospheric existence is crucial for comprehending the factors contributing to climate change and global warming [7, 19, 20].

Chapter - 3

Global Warming Trends in 2024

Air temperatures near the surface on Earth have been steadily increasing since the onset of the industrial age as a result of a combination of natural and anthropogenic processes. It is "extremely likely that more than half of the observed increase in the global average surface temperature from 1951 to 2010 can be attributed to the rise in greenhouse gas concentration due to human activities." The latest Intergovernmental Panel on Climate Change report, which was published in October 2013, underscored the significance of this issue by elevating the level of scientific caution with a remarkable 95% certainty in its main conclusions. The report provided detailed analyses and projections that indicated the overwhelming consensus among scientists regarding the rising temperatures and their link to human activities.

The global average temperature has experienced a notable rise of approximately 0.8 °C during the 20th century, with this trend notably accelerating over the past three decades. The scientific community has been closely monitoring this alarming increase in temperature, which has raised concerns about its consequences on the environment and human well-being. The rise in temperature has led to numerous repercussions, including melting ice caps, rising sea levels, more frequent and severe heatwaves, as well as changes in weather patterns.

It is important to highlight that nine of the hottest years of the last century can be traced back to the first decade of the 21st century. The data collected from various sources, such as weather stations, satellites, and oceanic measurements, provide compelling evidence of the escalating temperatures. This increase in temperature has far-reaching implications for ecosystems worldwide, causing shifts in plant and animal habitats and raising the risk of species extinction.

Notably, in 2012, the United States of America observed a new record of soaring temperatures, which stood as a striking example of the ongoing global warming phenomenon. The heatwaves that swept across the nation emphasized the urgency of adopting measures to mitigate climate change and reduce greenhouse gas emissions. The extreme weather events, such as hurricanes, droughts, and wildfires, have also become more frequent and intense, posing significant challenges for both human societies and natural ecosystems.

Moreover, the period from 2001 to 2010 emerged as the warmest decade in recorded history, particularly starting from 2005 onwards. This milestone further highlights the need for proactive measures to address climate change and its far-reaching impacts. World governments and international organizations have come together to formulate agreements and frameworks, such as the Paris Agreement, to tackle global warming and promote sustainable practices. These initiatives aim to reduce carbon emissions, increase reliance on renewable energy sources, and enhance the resilience of communities in the face of climatic changes.

It is crucial for individuals, communities, and nations to recognize the urgency and severity of the climate crisis and take collective action. By embracing sustainable lifestyles, transitioning towards clean energy, and advocating for policy changes, we can strive towards a more sustainable and resilient future. The expansion of renewable energy infrastructure, preservation of natural ecosystems, and adoption of climate-smart agriculture practices are just some of the steps that can contribute to mitigating the impacts of climate change.

In conclusion, the growing body of scientific evidence, combined with the observed trends in temperature rise, reinforces the need for immediate action to address climate change. The implications of global warming are far-reaching, affecting ecosystems, economies, and the well-being of present and future generations. By recognizing the urgency of this issue and acting collectively, we can work towards a more sustainable and livable planet for ourselves and future generations [21, 22, 23].

The global average UV radiation for this period was 0.5 W/m2 above the calculated long-term trend. This increase in UV radiation has significant implications for the environment and is closely linked to the rise in global average temperature. As the temperature continues to increase, various alarming phenomena have been observed, including a progressive rise in the sea level, the melting of Arctic and Antarctic ice, which combined contribute to increased flooding and changes in water patterns. Furthermore, there has been a severe drought in certain regions of the world, particularly in Australia, the Horn of East Africa, and parts of the Amazon Basin. These events serve as key indicators of the ongoing climate change crisis that the world is facing. It is important to note that the high concentration of atmospheric carbon dioxide (CO_2) has played a significant role in the cyclic temperature patterns observed over the past decade. The increase in CO_2 emissions has led to a greenhouse effect, trapping heat within the Earth's atmosphere and further amplifying the global warming process. This phenomenon has prompted researchers and scientists to develop reference scenarios in order to graphically represent the progression of global average temperature over time. These scenarios take into account both existing technologies and potential new technologies, while also considering regulatory strategies that can help stabilize carbon levels in the atmosphere. Among the most widely used reference scenarios are those presented by the Intergovernmental Panel on Climate Change (IPCC). These scenarios, known as A1B, A1T, B1, and B2, outline possible future trajectories based on different assumptions regarding economic, social, and technological developments. By utilizing these reference scenarios, scientists are able to assess the potential impacts of various policy decisions and technological advancements on global temperature trends. Ultimately, the goal is to find strategies and regulations that will effectively stabilize carbon levels in the atmosphere, mitigating the negative effects of climate change and ensuring a sustainable future for generations to come.

Due to these concerning observations and ongoing climate change, it is imperative to prioritize the adoption of sustainable practices and environmentally friendly innovations worldwide. This involves implementing policies and regulations that encourage the reduction of greenhouse gas emissions and the transition to alternative energy sources such as solar and wind power. Additionally, fostering international collaborations and agreements aimed at collectively addressing climate change is crucial. By joining forces, countries can share knowledge, resources, and technological advancements, leading to more effective solutions.

Moreover, understanding the complex interconnections between different ecosystems and their response to environmental changes is vital. This requires comprehensive scientific research and monitoring programs that provide valuable data on the impacts of climate change on habitats, biodiversity, and various species. By gaining deeper insights into these intricate relationships, policymakers and conservation organizations can make informed decisions to protect vulnerable ecosystems and safeguard the incredible diversity of life on our planet.

Recognizing that climate change affects not only the natural world but also human societies, it is essential to prioritize the development of resilience strategies. Building resilient communities includes investing in infrastructure that can withstand extreme weather events, developing early warning systems for natural disasters, and implementing sustainable agricultural practices to ensure food security. These measures can help minimize the social and economic impacts of climate change, particularly in vulnerable communities that are disproportionately affected.

Education and awareness play critical roles in combating climate change. Promoting environmental education from an early age and integrating climate change topics into school curricula can empower future generations to become environmentally conscious citizens. Additionally, support for public awareness campaigns and initiatives that inform individuals about the importance of reducing their carbon footprint and adopting sustainable lifestyles is crucial. Encouraging behavioral changes, such as conserving energy, reducing waste, and promoting eco-friendly transportation options, can make a significant collective impact.

Furthermore, climate skepticism is countered, addressing common skeptic claims and misrepresentations of climate science, some of which are even recognized among climate scientists as legitimate queries. These responses ultimately underscore the tenets of climate science, including the human influence on climate change, the future path of climate change given non-intervention, and the understanding of climate change by climate scientists. The basic tenets of climate science are deceptively simple, which may help to account for the 97% scientific consensus.

There is nothing wrong with doubt: it is a central component of the scientific method. Skepticism regarding climate change is distinct from legitimate scientific skepticism. Legitimate skepticism typically involves methodological concerns about climate measurements, models, or theoretical understanding, while general public skepticism disregards climate science altogether, with questions directed not so much at the scientific method but at the motives and honesty of climate scientists. Thus, outright rejection of, for instance, anthropogenic climate change is necessarily anti-scientific. Sources of controversy about climate change lie not in the science but in the politics; scientists, politicians, and the public are often necessarily ignorant regarding one another's interests and motivations. Climate policy is framed as a major intervention in the economy, raising issues of fairness and historical responsibility.

As the need for radical changes increases, an overwhelming amount of scientific evidence and popular opinion surrounding global warming has emerged. Generally, there are two positions toward global warming: those of concern and those of skepticism. The latter forms of argument are almost always metaphysical in nature and imperfectly consistent. By picking scientists' logical arguments apart, it is possible to greatly undermine the skeptical case. Picking cultural and investment-related arguments apart is much trickier and requires a good understanding of alternative worldviews. However, it is possible to argue that not only is it logically wrong to deny the possibility of anthropogenic global warming (AGW), but it is also morally irresponsible.

Basic arguments against concern are dismissed as stupid. Denying a causal connection between increasing carbon dioxide (CO_2) content in the atmosphere and increasing mean surface temperature is implausible in view of the long-range scientific evidence. It defies the laws of physical chemistry and well-known physical forces responsible and isotopically tracer aesthetically apparent globally and locally, and it is thus inconsistent with thousands of replicated and beautifully detailed instrumental and reconstructed records. Positive feedback effects are quantitatively well-documented. Forest cover and other biological ecosystems strongly impact the climate and are set to weaken or counteract the warming at high latitudes.

Skeptics argue that the warming stopped since 1998, and temperatures are well below the IPCC best-fit scenario prediction, which is utterly wrong. For avid skeptics, recent temperatures match previous warm epochs, and there is always a nearby volcano or an orbit effect. It is evident that a very simple energy-balance model comparing CO_2 with smaller radiatively active gases, such as methane and nitrous oxide, improves the fit mockingly well. Skeptics argue that current models do not account for internal variability or cloud feedback correctly and thus are untrustworthy. Such ignorance cannot be easily rebutted, but it is indicative of how far skepticism has strayed from science's healthy dilemmas.

Unlike weather forecasting, which typically addresses short-term projections, currently available climate change projections generally operate on longer timescales - years to decades. Efforts made through the IPCC/IS92 scenarios in the early 1990s spawned a huge array of climate change projections that were made using various general circulation models (GCMs) of the climate. These simulations provided some of the first estimates about how the projected climate change might vary regionally.

With the advent of the Special Report on Emissions Scenarios (SRES) scenarios, further climate change projections were generated. These, along with the IS92 climate change projections, were compiled into a dataset (the

SRES dataset) for public access. The SRES dataset of climate model experiments is widely applicable and is anticipated to be the principal resource for generating future climate impact studies at both the national and international scales.

Although there was independent replication of earlier IS92 climate change projections by different modeling groups, subsequent SRES climate change projections (with the exception of a handful of A1FI runs) have, to date, been made only by the PCMDI modeling groups. The SRES dataset remains an invaluable resource for assessing the potential impact of climate change on various national and international environmental issues, providing a common framework for policy-oriented scientific assessments involving climate change and impacts.

A common approach to examine the changes of climate using computer models is to describe scenarios with specified atmospheric CO_2 concentrations as a function of time. These scenarios can be calculated with climate models and by using different input greenhouse gas emission scenarios for CO_2. The desired concentration evolution can then be derived from the mass balance equation for CO_2 in the atmosphere. Mass balance calculations on CO_2 show that the atmospheric CO_2 concentration grows asymptotically towards an equilibrium concentration, depending on the respective CO_2 emission scenarios. For the selection of the emission scenarios, a wide range is available under various long-term constraints of CO_2 concentration stabilization and equilibrium temperature increase.

For the early industrial times, two scenarios are used as forcing. Because of its relevance for earlier discussions of climate scenarios, two 20th-century projections are applied in the first computations. Based on these atmospheric CO_2 concentrations over the past and current century, seven future climate scenarios are considered. Thus, over a time horizon of 100 years, GHG emissions will be strictly regulated on a constant decline to stabilize the CO_2 concentration in the atmosphere. Large-scale carbon sequestration has been applied, limiting the maximum allowable CO_2 concentration in the atmosphere to 550 ppmv or 720 ppmv. For projections using the A271 and B261 GHG emission scenarios, CO_2 concentrations will continue to rise rapidly beyond 2,100.

In climate scenario projections, all forcing is treated as constant externally imposed. Pre-industrial conditions enter the calculations as the initial value of the climate system, and terrestrial, oceanic, and cryospheric changes associated with climate warming have no dynamic feedback on the climate.

To account for uncertainties in the climate response to a given increase in CO_2 concentration levels, a range of model climate sensitivity (CS) parameters has been provided with climate evolution and sea level rise under different thresholds of CS parameter. Another important source of uncertainty is how different GHG emissions evolve. Each of the climate models can be forced with these four emission scenarios of GHG by anthropogenic activities. By using an integrated modeling framework, 21 long-term future climate and sea level scenarios are generated by coupling seven climate models and three GHG emissions scenarios.

Such a multi-model approach gives a broader probability distribution of the climate and corresponding sea level response to global warming than a scenario projected with one single climate model. The models are linearized with respect to both the forcing of GHG temperature rise in the atmosphere in response to a given increase in CO_2 concentrations and land-ocean heat capacity distribution.

It has become increasingly irrefutable that anthropogenic greenhouse gas emissions exert a profound influence on the climate system. Average global temperatures have increased in lockstep with rising concentrations of greenhouse gases in the atmosphere. The legitimacy of climate science has become a hotly debated topic among politicians and the public. Detractors and skeptics often point to past climatic conditions; alternatively, supporters espouse a view that forecasts near-term catastrophe. Scientists look at the best chances to mitigate the problem and, at worst, pragmatically hope to avert its most extreme long-term consequences.

Regardless of the differences, an extensive knowledge base has been amassed examining the coupling of surface temperature to simulations of future climate change resulting from anthropogenic perturbations to the atmosphere, a coupling found to be robust among models with different treatments of the atmosphere, ocean, sea ice, and land surface. Political action on global warming has so far remained reactionary, often inconsistent, and uneven across the globe; however, efforts to arrest the rapid and dramatic changes to Earth's climate will need to occur more vigorously in the near future, as delay will result in profoundly dire consequences. Without intervention, future generations may wish the Earth had not warmed.

Nevertheless, modeling efforts utilizing so-called earth-system approaches, often with explicit consideration of tipping points in the climate system, suggest that substantial and unpredictable changes may occur even during this century if greenhouse gas emissions are not sufficiently curbed.

Positive feedback mechanisms, such as the virtuous cycle of decreased albedo causing accelerated melt of polar ice sheets, may propel global temperatures into uncharted territory, with devastating global repercussions. The geography of life will be irrevocably altered, ocean circulation patterns may be disrupted, and the thermohaline circulation shut down, and methane hydrates may be released from tundra and permafrost regions. All seriously studied scenarios imply catastrophe.

The models also show that near-term interannual variability cannot simply be calculated using model-player perturbation pathways. Sudden unanticipated changes may occur due to chaotic dynamics. For example, the climate system could jump rather rapidly and quite unpredictably from one state of stability to another many degrees cooler, permanently reversing a trend of good weather that had prevailed for centuries or even millennia, and plunging regions with mild climates today into ice-age conditions. Rare events and "black swans" result from chaos when the timeframe is too short to realize the uncertainty. A probabilistic view of future abrupt change cannot be readily constructed with confidence. And as for the near- and long-term management of the climate system, freedom truly has uncertainties.

As the planet faces the dire consequences of climate change, global warming, and biodiversity loss, there is an urgent need for policymakers, world leaders, economists, businesses, and industries to come together to address these challenges. Climate policies, green new deals, and initiatives to combat this impending doom are important steps in the right direction. However, at the same time, each and every citizen has an equally important role to play as we all live on this one planet that bears no borders. Citizens have to understand and realize the importance of their individual and collective efforts to rectify and reverse some of the damage done.

Climate change, caused largely by greenhouse gas emissions, has altered weather patterns, heat waves, and rainfall distribution, resulting in a direct threat to food security, water supply, and extreme weather events such as mudslides, flash floods, cyclones, hurricanes, and forest fires. Sustainability has become crucial for continued societal advances, technological innovations, and economic growth. It is the responsibility of every inhabitant to care about this planet and ensure its sustenance. Awareness towards sustainability is equally important as it entails the adoption of economic, environmental, and social practices that will benefit all living beings and the planet. It is crucial for a healthy earth as well as a healthy economy and society.

However, sustainability has gained popularity only in the recent past. The ongoing efforts towards sustainability have not yet been materialized to the fullest and there is a need to identify the impediments in its path. Studies have shown that there is awareness about the concept of sustainability among people, but at the same time, preference towards the protection of the immediate surrounding, family, and community has also been indicated. This indicates a lack of information towards awareness of larger problems such as climate change and poverty and at the same time highlights the challenges of a broader global mindset.

Addressing climate change requires a multilateral approach that involves not only governments but also businesses, civil society, and individuals. By fostering collaboration and shared responsibility, it is possible to create a sustainable future for our planet. Through technological advancements, policy innovations, and concerted efforts, we can mitigate the negative effects of climate change and pave the way for a thriving and resilient world for current and future generations. It is our collective responsibility to take action now and ensure a healthier, more sustainable planet for all [24, 25, 26].

3.1 Temperature changes

The latest global temperature of land, recorded from January to October in the year 2024, was determined to be 0.47°C higher than the average temperature observed between 1981 and 2010. This surpasses the previous record set in 2023, making it the warmest period on land since the year 1875. Several factors contributed to this significant increase in temperature, aside from the ongoing trend of year-by-year warming. These factors include a weak La Niña event, which resulted in cooler temperatures during the years 2022 and 2023, as well as a substantial year-to-year increase in warmth, ultimately leading to the exceptionally high temperatures in 2024. The findings from this study further corroborate the accelerating trend of global mean air temperature witnessed since the commencement of the 21st century. Moreover, the recorded tropospheric temperature for the year 2024 demonstrates a noteworthy 0.74°C increase above the long-term average observed in 1875. The Tropic region, specifically, experiences the most significant rise in temperature, reaching an alarming 1.01°C above the average. These rising temperatures have profound implications for various geographic regions. For instance, the subtropical East and Western Hemisphere oceans, along with the higher latitudes in the Northern Hemisphere, witness substantial increases in not only heat content but also air moisture. These findings suggest that the year 2024 has further amplified the transfer of heat and moisture from the upper portions of the ocean into the atmosphere, subsequently reducing the likelihood of near-future cooling. Given the continuous concentration of atmospheric greenhouse gas forcing, there is a growing risk associated with the global climate during the summer season. The persistent accumulation of greenhouse gases in the atmosphere intensifies the greenhouse effect, leading to elevated temperatures and potentially adverse climatic conditions worldwide. It is imperative that effective measures are implemented to address this concerning issue and mitigate the potential consequences of climate change on a global scale.

The consequences of these rising temperatures are multifaceted and far-reaching. One of the most apparent impacts is the accelerated melting of the polar ice caps, which contributes to rising sea levels and threatens coastal communities. Additionally, the warming trend has significant implications for the world's ecosystems. As temperatures continue to rise, many plant and animal species struggle to adapt to their changing environments, leading to disruptions in biodiversity and potential extinctions. Furthermore, extreme weather events such as hurricanes, droughts, and heatwaves are becoming more frequent and intense, causing immense societal and economic damage. These events not only endanger lives but also pose significant challenges to agriculture, water resources, and infrastructure.

The year 2024 serves as a clarion call for immediate action to combat climate change. It is crucial that governments, industries, and individuals prioritize sustainable practices and reduce greenhouse gas emissions. Transitioning to renewable energy sources, such as solar and wind power, can help mitigate the environmental impact of fossil fuels. Additionally, efforts to conserve and protect natural habitats are essential for preserving biodiversity and ensuring the long-term health of ecosystems. Implementing policies that promote energy efficiency, sustainable agriculture, and responsible consumption can also contribute to climate change mitigation.

Addressing climate change requires global cooperation and commitment. International agreements, such as the Paris Agreement, provide a framework for countries to collaborate in reducing emissions and adapting to the effects of climate change. Continued investment in climate science and research is vital to expanding our understanding of the Earth's climate system and informing sound policy decisions. Education and public awareness campaigns are also essential to fostering a collective sense of responsibility and encouraging sustainable lifestyle choices.

The year 2024 has highlighted the urgency of the climate crisis and the need for immediate action. If we fail to curb greenhouse gas emissions and mitigate the impacts of climate change, future generations will bear the brunt of our inaction. By taking decisive steps now, we can create a sustainable and resilient future for ourselves and the planet. Let us seize this opportunity to address climate change head-on and pave the way for a healthier and more sustainable world [27, 28, 29].

Impacts: The 2024 temperature was 0.47°C above the 1981-2010 average, surpassing the record set in 2023 to become the warmest year on record since 1875. The unprecedented warmth of 2024 not only confirmed the ongoing trend of rising temperatures but also highlighted the urgency of addressing the underlying causes of climate change. Over the past few decades, the world has witnessed a steady increase in global temperatures, with 2024 marking a significant milestone in this relentless upward trajectory. The period between 2000 and 2024 has proven to be particularly alarming, as the average temperature during this time has consistently remained well above the levels recorded in the latter half of the 20th century. With a staggering trend of 0.24°C per decade, the rate of temperature increase has reached its highest point since the start of instrumental measurements in 1875. This clear evidence of continuous warming calls for immediate action to curb greenhouse gas emissions and mitigate the devastating effects of climate change.

It is crucial to consider the factors that have contributed to the notable temperature variations within the past decade. While the years 2022 and 2023 experienced relatively cooler temperatures, the overall trend remained steadfast, owing to a combination of variables including a slight La Niña, a predominantly cool period between 2022 and 2023, and a consistent upward trajectory from one year to the next. Although the rate of temperature increase in 2023 was greater than that of 2024, the overarching seven-year trend clearly indicates an acceleration in global mean air temperature, providing further evidence of the escalating warming trend in the 21st-century. The pressing question remains: Are these unprecedented temperature increases a direct consequence of continuous anthropogenic greenhouse gas emissions? The scientific consensus overwhelmingly points to human activities, namely the release of greenhouse gases into the atmosphere, as the primary driver of global warming. The burning of fossil fuels, deforestation, and industrial processes have all contributed to an unprecedented accumulation of greenhouse gases, trapping heat within the Earth's atmosphere and causing the planet to warm at an alarming rate.

Unless immediate and decisive actions are taken to reduce emissions and transition to cleaner, more sustainable energy sources, the current trajectory of accelerating warming could have catastrophic consequences for our ecosystems, economies, and the well-being of future generations. It is our collective responsibility to confront this crisis and work towards a sustainable and resilient future for all. The year 2024 serves as undeniable evidence of the progressive climate crisis we are facing. With temperatures 0.47°C above the 1981-2010 average, surpassing the previous record set in 2023, it has become the warmest year since records began in 1875. This alarming heat not only strengthens the ever-present trend of rising temperatures but also emphasizes the critical need to address the root causes of climate change.

Throughout the past decades, global temperatures have steadily risen, and 2024 is a major milestone in this alarming ascent. The period from 2000 to 2024 has been particularly worrisome, with average temperatures consistently exceeding those recorded during the latter half of the 20th century. The rate of increase, a staggering 0.24°C per decade, reaches its highest point ever since instrumental measurements began 149 years ago. The undeniable proof of this continuous warming demands immediate action to counteract greenhouse gas emissions and mitigate the catastrophic effects of climate change. We must carefully analyze the factors contributing to the notable temperature fluctuations within the past decade. Despite relatively cooler years in 2022 and 2023, the overall trend remains resolute due to various factors, including a mild La Niña, a predominantly cool phase from 2022 to 2023, and a steadfast upward trajectory year after year. Although the rate of temperature increase in 2023 surpassed that of 2024, the comprehensive seven-year trend clearly indicates an acceleration in global mean air temperature and further supports the undeniable escalation of warming in the 21st century.

The critical question that arises is: Are these unprecedented temperature increases a direct consequence of continuous anthropogenic greenhouse gas emissions? Scientific consensus overwhelmingly attributes these changes to human activities, specifically the release of greenhouse gases into the atmosphere, which act as the primary catalyst for global warming. The combustion of fossil fuels, deforestation, and industrial processes have collectively contributed to an unparalleled accumulation of greenhouse gases, resulting in the entrapment of heat within the Earth's atmosphere and an alarming rate of planetary warming. Unless immediate and resolute actions are taken to reduce emissions and transition to cleaner and more sustainable energy sources, the current trajectory of accelerated warming could yield catastrophic outcomes for our ecosystems, economies, and the future well-being of succeeding generations. It is our shared responsibility to confront this crisis head-on and strive diligently towards a sustainable and resilient future for all [30, 31, 32].

Chapter - 4

Impact on Weather Patterns

As global temperatures continue to rise at an alarming rate, the consequences for weather patterns are set to become even more pronounced. The intensely powerful tropical storms we currently witness will unleash their fury with unprecedented force and ferocity in the coming years. The havoc they wreak on our environments and ecosystems will be unimaginable, leaving a lasting impact on the crops we rely on for sustenance.

The cultivation of crops such as corn, wheat, and soybeans, long considered staples of our diets, will be profoundly affected by these looming changes. These plants have evolved to thrive under specific climatic conditions, carefully embracing the perfect blend of heat, moisture, and sunshine. Yet, this delicate balance, painstakingly achieved through centuries of agricultural innovation, is now under threat. The smallest alteration in the availability of these vital components can disrupt their growth, causing a significant reduction in yields and subsequently impacting the cost of our food.

However, the challenges do not end there. With just a few degrees increase in temperature, the resulting evaporation will intensify, leading to the displacement of crops from their once-optimal environments. What was once a fertile land for corn may become arid and unsuitable, forcing farmers to explore new territories and adapt their practices. Similarly, excessive or insufficient rainfall may plague other crops, disrupting their delicate equilibrium and leaving them vulnerable to disease and stunted growth. The consequences on our food supply are dire.

Furthermore, plants exhibit their peculiarities when it comes to nutrient intake, much like selective eaters in the animal kingdom. Understanding these intricate preferences becomes paramount as we navigate the challenges posed by shifting weather patterns. The careful management of nutrients, ensuring that each plant receives its required nourishment, will be essential in maintaining healthy and productive crops.

Therefore, it is imperative that we not only acknowledge the impending changes but also take proactive measures to safeguard our agricultural practices and protect our food supply. Adapting to evolving weather patterns demands a comprehensive approach, involving cutting-edge technology, innovative farming techniques, and a deep understanding of the intricate factors at play. By embracing these challenges head-on, we can ensure a resilient and sustainable food system that withstands the test of time [33, 34, 35].

Over the course of the past century, spanning a substantial 100 years, the average global temperatures across the planet Earth have undergone a significant and noteworthy increase, precisely amounting to a rise of approximately 1 degree Celsius. Numerous esteemed researchers and experts within the scientific community vociferously contend that the phenomenon commonly referred to as global warming possesses the potential to instigate an assortment of even more extreme weather conditions, ultimately causing climate change to accelerate at an unprecedented pace. In fact, it is contended that within the impending 50 to 100 years, various regions across the globe may bear witness to the exacerbation of heat waves in terms of both frequency and severity. Such a foreboding scenario can be comprehensively attributed to the incremental rise in the levels of carbon dioxide (CO_2) present within the Earth's atmosphere. The very escalation of these detrimental climatic phenomena, such as severe storms, intense heatwaves, and copious amounts of heavy precipitation, have become unequivocally intertwined with the aforementioned surge in CO_2 levels. Consequently, this unrelenting surge in heat shall inevitably precipitate elongated periods of debilitating droughts, accompanied by a commensurate augmentation in the atmospheric water vapor load. Intriguingly, this exacerbation in the volume of atmospheric moisture accentuates the intensity and frequency of heat waves and the consequential heavy precipitation, largely attributable to the heightened process of evaporation. It is of paramount importance to comprehend that the inextricable link between global warming and climate change vis-à-vis weather patterns and weather-related events is of a most significant nature. Such a nexus indubitably precipitates a direct impact on various facets of our daily activities, exerting substantial influence over an array of tasks both mundane and critical in nature. These tasks commonly include our occupational endeavors, recreational pursuits, journeys and voyages, and even the most banal of errands that punctuate our quotidian lives [36, 37, 38].

4.1 Extreme weather events

Over the past five decades, there has been a significant and deeply concerning surge in the frequency and magnitude of extreme weather events occurring across the globe. These catastrophic and devastating events, encompassing scorching heat waves that scorch the earth's surface, wreak havoc on delicate ecosystems, and pose a grave danger to human health. In addition to the blistering heat waves, we witness the devastating power of hurricanes as they increase in intensity and unleash their destructive force, leaving countless regions in ruins. Moreover, severe tropical storms and relentless blizzards pound vulnerable areas with merciless fury, burying cities and towns under heaps of snow and ice. The rising global temperatures also contribute to unprecedented and catastrophic flooding, submerging once-thriving communities and displacing entire populations. The toll of these extreme weather events is immeasurable, as they bring widespread devastation, loss of life, and irreversible damage to our natural world.

If the current trends persist, leading experts confidently forecast that by the year 2024, the reverberating impacts of global warming will transform these extreme weather events into a terrifyingly commonplace occurrence. The world as we know it will be forever altered, and our ability to adapt and withstand these climatic challenges will be put to the test. There is no doubt that urgent action is required to mitigate the dire consequences that await us if we fail to address this crisis swiftly and effectively.

One of the most disconcerting ramifications of the alarming heat waves exacerbated by global warming is the peril they pose to human health. As temperatures soar to unprecedented heights, a myriad of major cities across the globe have already borne witness to a calamitous surge in fatalities, traffic accidents, and hospital admissions. On the sweltering days of intensifying heat, these already bustling cities become tragic witnesses to a doubling, and in some cases even tripling, of these unfortunate incidents. The toll on vulnerable populations, such as the elderly, children, and individuals with pre-existing health conditions, is particularly devastating. In actuality, prolonged exposure to extreme heat has the potential to induce severe illnesses, including heatstroke, dehydration, and cardiovascular complications, and tragically, even fatalities. The healthcare systems in these affected regions are pushed to their limits, struggling to cope with the influx of patients and grappling with the shortage of resources and facilities.

Furthermore, the mounting sea levels precipitated by global warming pose an imminent and formidable threat to low-lying regions worldwide. Let us contemplate the city of Seattle in the state of Washington, for instance. As the inexorable rise in sea levels continues unabated, the risks of destructive flooding loom ominously over this vibrant metropolis, once known for its stunning coastal landscapes. The picturesque waterfront neighborhoods, bustling with life and commerce, now face an uncertain and perilous future. In anticipation of this environmental hazard, metropolises across the globe, including Seattle, have already begun implementing extensive modifications to their city streets. These modifications encompass a broad range of measures, including the necessary widening of streets to accommodate the increased volume of water flow, thus safeguarding against potential damages of cataclysmic proportions. These adaptations are crucial in the face of this looming threat, as they strive to protect lives, properties, and the overall well-being of the affected communities.

Without a shred of doubt, the impacts of global warming on extreme weather events are undeniably poised to reshape our world in manners we simply cannot afford to overlook. The urgency to promptly address and mitigate these looming consequences cannot be overstated. Any failure to execute timely and decisive actions may result in irreparable damage to the delicate ecological systems, communities, and economies that underpin our societies. It is of paramount importance that governments, organizations, and individuals unite in a concerted effort to implement substantial and sustainable solutions to combat global warming and the consequential escalation of extreme weather events. Only through the strength of collective action can we dare to hope for a safer and more resilient future, one that embraces the well-being and prosperity of all [39, 40, 41, 42, 39, 40, 43].

As the global climate continues to participate in global warming, it's not out of the ordinary that we are going to see a significant increase in other windy seasons. The intensified windier seasons create a greater number of hurricanes and tropical storms that are significantly more intense compared to previous years. In fact, the severity of these storms can be overwhelming. Additionally, the impact of these storms can reach uncharted territories as hurricanes are projected to occur even further north than usual. By the year 2024, the Northern Hemisphere echo-tropics will experience a remarkable shift towards tropical storms and hurricanes. This shift will result in a drastic increase in stormy weather characterized by extreme conditions. These weather phenomena will take place predominantly along the notorious Hurricane Alley, which will gradually shift further north than what has been witnessed in previous years. Consequently, this alteration will contribute to a surge in flooding incidents particularly in the southern and southeastern parts of the nation. The increased frequency and intensity of storms will result in heightened risks of widespread flooding and its detrimental aftermath. Moreover, the change in climate patterns will also have a profound impact on coastal regions, leading to erosion, loss of habitats, and threats to marine ecosystems. Furthermore, during the winter season, the Northern Hemisphere will bear the burden of amplified blizzards and snowstorms. The intensification of these winter weather events will affect not only the usual regions but also extend their reach to southern cities that are not adequately equipped to deal with such extreme weather conditions. These cities, unaccustomed to heavy snowfall, may encounter significant challenges in managing the consequences of such storms on their infrastructure and daily activities. This unexpected occurrence of severe winter storms in areas unaccustomed to such events may lead to disruptions and hardships for the affected communities. In addition to the immediate and localized impact of these weather events, there are broader ecological and environmental implications. The increased frequency and intensity of storms can result in significant damage to natural ecosystems, including deforestation, loss of biodiversity, and disruption of delicate ecological balance. Coastal areas, in particular, face the risk of saltwater intrusion, which can harm freshwater sources and impact agriculture. Furthermore, the rise in temperature due to global warming has the potential to lead to the melting of ice caps and glaciers, contributing to rising sea levels and posing a significant threat to low-lying coastal regions. These regions will not only be susceptible to severe weather events but also face the risk of displacement and loss of livelihoods for their inhabitants. In conclusion, the continuous progression of global warming is resulting in a series of intricate consequences on our climate. The anticipated increase in windy seasons is contributing to a surge in the number and intensity of tropical storms and hurricanes. Moreover, the altered patterns are causing these weather phenomena to occur in regions further north, resulting in the need for adaptation and preparedness. The effects of these storms are not limited to the immediate areas but have far-reaching consequences, such as increased flooding in the southern and southeastern regions. Additionally, the amplified winter storms pose a daunting challenge for southern cities that lack the necessary infrastructure to cope with such events. As we approach the year 2024 and beyond, it becomes imperative to address these changes in climate patterns and take proactive measures to mitigate their adverse impact on the human population, wildlife, and the environment at large. Only by understanding the gravity of the situation and working together can we ensure a sustainable future for all. It is crucial for countries to cooperate and implement policies that reduce carbon emissions and transition to renewable energy sources. Furthermore, enhancing coastal defenses and investing in resilient infrastructure can help protect vulnerable communities from the increasing risks posed by extreme weather events. Educating the public and raising awareness about the importance of climate change adaptation and mitigation strategies is also essential. By promoting sustainable practices and reducing our ecological footprint, we can minimize the negative effects of global warming and create a more resilient and sustainable future. It is our collective responsibility to safeguard the planet for future generations, and by taking decisive action today, we can make a significant difference in shaping a better tomorrow. Let us unite and prioritize the preservation of our environment, for it is our duty to leave behind a flourishing planet that sustains life and brings harmony to all living beings. The need of the hour is to embrace renewable energy sources, innovate greener technologies, and foster a sustainable mindset. By fostering international collaborations, instituting stringent environmental regulations, and instilling eco-consciousness in every aspect of our lives, we can combat the adverse effects of climate change and build a thriving future for generations to come. Together, let us embark on this journey towards a greener, cleaner, and more vibrant world, where the profound beauty of nature thrives and sustains humanity.

Chapter - 5

Impact on Ecosystems

Ecosystems of cities and regions. With the rapid growth of the urban population, ecosystems are changing under the influence of intensified environmental pollution. According to forecasts, in 2024 their state will be very tense, as the consequences of man-made accidents and the problems of water storage, air and soil composition will become more acute. Pending environmental problems of large cities will be related to maintaining urban ecosystems that will provide the population of megacities with food, oxygen, green and water tourism. Also, indoor green spaces in the form of living walls, green tubes installed in residential buildings, and the construction of vertical farms are becoming popular. Indoor urban ecosystems include rooftop gardens, balconies and urban farms. Scientists predict that vegetation on the facades of buildings in the high commercial centers of large cities will expand, which will provide life-sustaining space and species diversity for insects, other small organisms and bird populations. The development of these urban ecosystems will make it possible to reduce the negative environmental impact in the future, both of man-made objects and the activities of the residents.

5.1 Loss of biodiversity

The concept of endangered species and the loss of biodiversity are not new concepts to the human race but often misunderstood. The idea is that if a species or a group of species is completely wiped off the Earth through an extinction process, it can never be brought back ever again. What is lesser known is the danger and negative impact to the human race as the result of the loss of biodiversity. Genetically diverse species, ecosystems, and niches provide the human race with the air we need to breathe, the food we need to eat, and the environment we need to live in. By losing any of these vital species and ecosystems, the human race will suffer and struggle to survive. The loss of biodiversity, therefore, is not only a vital concern but also another indirect piece of evidence indicating that the Earth is currently undergoing a new mass extinction, one that is primarily caused by the actions of the human race.

To date, we have only discovered one natural evolutionary process to bring back extinct species and ecosystems. It's called Evolutionary Time.

Unfortunately, this super slow process will take many millions of years to occur and is completely beyond anyone's control. The loss of biodiversity in the modern world is influenced by a combination of profoundly intricate and multifaceted factors in both natural and human-induced situations. In the wondrous realm of nature, exquisite animals and vibrant plants tirelessly engage in fierce competition with one another for the priceless bounty of limited resources nestled within their intricate ecosystems and unique niches – resources that encompass not only sustenance, but also sanctuaries, compatible mates, and immaculate breeding grounds. A myriad of influential forces, including but not limited to disease, predation, parasitism, fierce competition, and unyielding environmental transformations, relentlessly act as the guardians that regulate the population size and behavioral patterns of the diverse array of species inhabiting the wild [44, 45, 46].

Chapter - 6

Impact on Human Health

Global warming is now thought to have exerted an influence on the human population worldwide through changes in the climate factors. In the year 2024, mass media have frequently reported on heat and cold injuries occurring during extreme weather, as well as on infectious diseases such as dengue fever, malaria, barreliasis, encephalitis, chikungunya, Nile virus, epidemic polyarthritis, Lyme disease, and cholera being increasingly found in temperate zones. The geographical changes in the habitats of the nematode Baylisascaris and the worm Schistosoma affecting domestic animals have also been reported by epidemiologists. Crops, fruit trees, vegetables, and ornamental plants have all responded with changes to the increase in the level of atmospheric carbon dioxide and other gases, to temperature increases, to increases in levels of rainfall, ultraviolet rays, atmospheric dust, and atmospheric pollutants, as well as to changes in relative humidity, resulting in, for example, decreased levels of nutrients in lettuce and spinach.

The production of poisonous substances causing food poisoning or component degradation in foods such as apple juice, olive oil, walnuts, horsetail, raw milk, tributyrin, shellfish, chichi, damanj, damaku, bamboo shoots, remembered beans, cashews, and figs, or even causing the production of biologically toxic substances has given rise to increasing consumer concern. This concern extends not only to the aforementioned food items, but also extends to various other categories including seafood, processed meats, dairy products, fruits, vegetables, and bakery items. The alarming rise in cases of food contamination has raised questions about the safety measures implemented in the food production industry. One particular example of this alarming trend can be observed in the context of the meat industry, especially the production of Japanese beef. Japanese beef, known for its premium quality and high demand in international markets, has not been immune to the concerns regarding consumer safety. Consequently, stringent regulations and protocols have been implemented to ensure the safety of this product. These regulations include the monitoring of cattle farms, regular health screenings of the animals, and strict adherence to hygiene practices in slaughterhouses and processing facilities. The issue of consumer safety and health extends beyond food production and encompasses various other aspects of modern lifestyles. In the field of vocational health, concerns have been raised about the excessive use of drugs and medications such as powerful antibiotics, first-generation antimicrobials, antifungals, demeclocycline, and cyclosporin of plant origin. These drugs, although intended to treat various ailments and infections, have inadvertently exposed individuals to an increased risk of oxidants and electrosmog. This phenomenon has sparked discussions and debates within the medical community regarding the potential long-term effects of these substances on human health. As consumers become more informed and conscious of the potential risks associated with food contamination and exposure to harmful substances, there is a growing demand for transparent labeling practices and stricter regulations in industries related to food and healthcare. Governments and regulatory bodies around the world are working towards implementing more robust systems for monitoring and ensuring the safety of food products and medications. In addition, consumers are encouraged to educate themselves about the potential risks and make informed choices when it comes to purchasing and consuming various products. Overall, the increasing concerns about the production of poisonous substances, food contamination, and exposure to harmful substances highlight the importance of consumer awareness, industry regulations, and proactive measures to safeguard public health. It is crucial for all stakeholders, including governments, businesses, and individuals, to collaborate and prioritize the well-being of consumers in order to mitigate these risks and ensure a safe and healthy future for all. In light of these issues, it is essential for authorities to invest in research and development for innovative technologies and strategies that can effectively detect and eliminate toxic substances from the food supply chain. This will not only enhance consumer safety but also bolster trust in the food industry. Furthermore, it is imperative for governments and regulatory bodies to establish comprehensive and standardized testing procedures to continuously monitor food products for contaminants. By implementing stringent protocols, more effective quality control measures can be put in place to prevent the occurrence of foodborne illnesses and protect the public from harm. Moreover, the development and enforcement of stricter regulations would act as a deterrent to unscrupulous individuals and corporations seeking to cut corners and compromise the safety of consumers. In addition to regulatory measures, it is essential for consumers to take an active role in safeguarding their own health. By being vigilant and informed, individuals can make better choices when it comes to purchasing and consuming food products. This includes reading labels, understanding ingredient lists, and seeking out products that are certified as safe and free from harmful substances. Additionally, consumers should prioritize buying from reputable sources and supporting businesses that prioritize consumer safety and wellbeing. To foster greater transparency and accountability, companies should consider implementing comprehensive labeling practices that not only disclose the ingredients used but also provide information on the sourcing, processing, and testing of their products. This would enable consumers to make informed decisions based on accurate and reliable information. Furthermore, companies should actively engage in research and development to identify and implement innovative technologies and techniques that reduce the risk of contamination and ensure the production of safe and healthy food products. The role of education and awareness cannot be understated in addressing the concerns surrounding food safety and consumer health. Governments, educational institutions, and healthcare providers should collaborate to develop educational programs aimed at raising awareness about the potential risks and promoting responsible consumption practices. By equipping individuals with knowledge and empowering them to make informed choices, the likelihood of foodborne illnesses and exposure to harmful substances can be significantly reduced. In conclusion, the increasing concerns about the production of poisonous substances, food contamination, and exposure to harmful substances have led to a growing focus on consumer safety and well-being. It is necessary for all stakeholders, including governments, businesses, and individuals, to prioritize the implementation of stringent regulations, transparent labeling practices, and proactive measures to mitigate these risks. By fostering collaboration and accountability, it is possible to ensure a safe and healthy future for all, where consumers can have confidence in the products they purchase and consume [47, 48, 49].

6.1 Heat-related illnesses

Heat-related illnesses like heat stroke, heat exhaustion, and dehydration can all lead to serious health effects, including heat stress and death. Children, the elderly, people who are overweight, and those with serious medical conditions are at an increased risk of heat-related illness. In the United States and other parts of the world, heat waves are becoming an increasingly serious public health threat. In fact, excessive heat events can cause more annual deaths than hurricanes, lightning, tornadoes, floods, and earthquakes combined. Increases in global temperature mean increases in heat waves, and some of these heat waves are expected to last for longer periods of time. These changes in heat will likely lead to more frequent and longer-duration heat waves, which is a growing public health problem.

Heat waves also place additional strain on the infrastructure of communities, including their health and utility systems. The increased temperatures can lead to a surge in demand for medical services, overwhelming medical facilities and necessitating the establishment of cooling centers and shelters. In buildings lacking air conditioning, the heat-sensitive load escalates, putting occupants at risk. In densely populated urban areas experiencing prolonged and widespread heat waves, there is a heightened possibility of brownouts and even blackouts, coinciding with the critical period when air conditioning is indispensable for survival. If the duration of the peak heat extends and the temperatures rise significantly, the overall heat within structures can accumulate to perilous levels, posing a threat to human life and well-being. Additionally, the impact of heat waves extends beyond the immediate health risks and infrastructure strain. Heat waves have far-reaching economic consequences, affecting various sectors and industries. For instance, agricultural productivity takes a hit as crops and livestock suffer from extreme heat stress, leading to lower yields and potential losses for farmers. The energy sector also faces challenges during heat waves, as power demands skyrocket due to increased cooling needs, putting pressure on power generation and distribution systems. This can result in higher energy costs for consumers and potential disruptions in the electricity supply. Furthermore, heat waves exacerbate social inequalities and disparities, disproportionately affecting vulnerable populations. The elderly, children, individuals with pre-existing medical conditions, and those living in poverty are particularly susceptible to the adverse effects of extreme heat. Lack of access to proper cooling systems, limited financial resources, and inadequate infrastructure in marginalized communities can amplify the health risks and hardships faced during heat waves. The social fabric of communities can also be strained, as individuals and families struggle to cope with the physical and emotional toll of prolonged heat exposure. Additionally, schools may need to close during extreme heat, disrupting the education of students. Addressing the challenges posed by heat waves requires a comprehensive approach that encompasses both short-term emergency measures and long-term climate resilience strategies. Public health agencies and emergency responders play a crucial role in disseminating heat warnings, providing support to vulnerable populations, and ensuring access to cooling centers and shelters. Urban planning and infrastructure development should prioritize the implementation of heat-mitigating measures, such as green spaces, cool roofs, and reflective pavements, to reduce the urban heat island effect. HVAC systems and building codes should be updated to include heat wave resilience standards. Climate change mitigation efforts, including reducing greenhouse gas emissions and transitioning to clean energy sources, are also essential in preventing the escalation of heat waves and their adverse impacts. In conclusion, the consequences of heat waves extend far beyond the immediate health risks, impacting infrastructure, economies, and social dynamics. As extreme heat events become more frequent and intense due to climate change, it is crucial to prioritize adaptation and resilience strategies to protect communities from the growing threat of heat waves. By addressing the multifaceted challenges associated with heat waves, societies can enhance their capacity to cope with and minimize the impacts of these extreme weather events. Building community partnerships and fostering public awareness and preparedness are vital for effective heat wave response and protection. Investing in research and development to improve heat wave forecasting and climate modeling can also contribute to better planning and mitigation efforts. Ultimately, a collaborative and proactive approach is required to build resilience and mitigate the impacts of heat waves, ensuring the well-being and safety of communities now and in the future [50, 51, 52].

Chapter - 7

Mitigation Strategies

Given the current knowledge on the causes of global warming and climate change, particularly the role of pollutants such as CO_2, we can go a step further than with other forms of pollution. Firstly, adopting clear and ambitious political objectives and fixing a time horizon for making the decisions and commitments required are of the essence. Secondly, we need to do some serious research. We do not know how much CO_2 we can produce before these emissions cause the harmful warming or climate change that we want to avoid. Once we are sure of the level of pollution that can be allowed, this allows businesses to engage in longer-term planning. It is vital that, as part of efforts to protect the environment and achieve sustainable development, governments, businesses, scientists and non-governmental organizations (NGOs) work together on mitigating climate change, and draw up a set of policies and strategies.

Governments must play a decisive role, setting a clear regulatory framework, fixing time horizons and implementing measures to achieve change in the longer term for public infrastructures. The private sector also has a significant role to play in order to achieve sustainable development. Businesses need not only to adopt responsible management practices but also to seek out transparent, credible business partners and suppliers that can be trusted. In the case of decisions that involve citizens, it is often questioning by civil society that decides the choices made. Information and dialogue are the main tools for involving civil society in this process. Finally, a system of fair and equitable sharing of both the costs and the benefits must be found, promoting relations built on trust and solidarity.

7.1 Renewable energy sources

Climate change is occurring at an extremely alarming rate and it is abundantly clear that humans are the primary culprits responsible for this dire situation. Among the numerous industries that contribute to this devastating phenomenon, the food and beverages sector emerges as one of the most nefarious polluters on a global scale. The pivotal question that arises is: How can we counteract this prevailing issue? The answer lies within a growing segment of the population that is fervently advocating for the adoption of a sustainable diet.

Across the globe, meat stands as the most extensively consumed food item. However, the production and consumption of meat generates an exorbitant amount of carbon dioxide emissions per kilogram, thus exerting an unmistakably detrimental impact on our climate. Nevertheless, it is important to note that humans cannot simply refrain from eating. Therefore, it becomes crucial to explore alternative solutions that can alleviate the strain we place on the environment.

In this battle against global warming, renewable energy sources emerge as indispensable allies. By harnessing these sustainable energy sources, we can effectively enhance energy efficiency and enable the utilization of a diverse range of resources. Whether it be solar power, wind energy, hydropower, or geothermal sources, the untapped potential of renewable energy is vast and holds the key to mitigating climate change. Not only do these sources provide a greener alternative, but they also possess the capacity to drive economic growth, create jobs, and foster technological advancements.

By embracing renewable energy, we can effectively combat the threatening repercussions of climate change. It is imperative that we acknowledge the gravity of the situation and take action now, for the preservation of our planet and the well-being of future generations. The power to catalyze change lies within our hands, and it is essential that we fervently advocate for the sustainable practices that will ultimately shape the future of our world [53, 54, 55].

Renewable energy sources benefit the environment since they, through the development of cutting-edge solar collectors or panels, advanced photovoltaic systems, and innovative wind technology, effectively prevent harmful air and soil pollution and, in most cases, conserve precious land resources. Given the undeniable reality of climate change, the adoption of these sustainable technologies is not only crucial but also an urgent ecological imperative. It has been more than a century since scientists first studied the greenhouse effect and global warming and warned that burning fossil fuels, oil, coal, natural gas, and other activities have increased the levels of industrial gases in the atmosphere. Global warming has already raised global temperatures and caused climate change, but the change, which is now locked in for centuries or millennia, is still small on top of natural changes over geological timescales. Only recently, dissent or questioning of climate science became a mass phenomenon, especially in the United States.

Global warming refers to the increase in the average temperature of the Earth's atmosphere due to an increase in greenhouse gases, principally carbon dioxide (CO_2) concentration. The greenhouse gases include CO_2, methane (CH_4), nitrous oxide (N_2O), chlorofluorocarbons (CFCs), and water vapor. Two of these greenhouse gases, CO_2 and CH_4, together account for more than 67 percent of the current greenhouse effect, with CO_2 alone accounting for about 56 percent. In addition to these gases, there is an abundance of solid and liquid pollutants like black carbon (soot) and urban-industrial haze (smoke and wet sulfuric acid) that affect the thermal balance of the Earth.

Pollution of the atmosphere with black carbon (soot) is a significant problem in many places (e.g., the US East Coast, Canada, and especially the Arctic), and soot is thought to contribute to glacial melting in several places (and thus indirectly contribute to sea level rise). However, soot is not a greenhouse gas, and there is an enormous difference between a solid (like soot) or liquid pollutant that scatters and absorbs various wavelengths of radiation and a gas pollutant (like CO_2) that only absorbs and re-emits long infrared wavelengths.

Global warming is expected to continue due to a combination of delayed responses in the climate system and continuing human emissions of greenhouse gases. Because of its long-term climate stability, the present sea level under today's climate, which is evidently ice-covered Antarctica, would not exist without natural greenhouse gas regulation. Major problems facing mankind with regard to climate change are highlighted here, but because of the scale of the problem, only a rather simple treatment can be offered.

According to the United States National Aeronautics and Space Administration (NASA), global warming or climate change is defined as "a long-term change in the average weather patterns on Earth". Global warming refers to the persistent rise in average surface temperature of the Earth's atmosphere, oceans, and landmasses. The greenhouse effect is a natural phenomenon where certain gases in the atmosphere trap heat and radiate it back to the Earth's surface. The Earth's temperature is determined by balancing the amount of energy it receives from the Sun and the energy it radiates back to outer space. The excess amount of radiated energy is absorbed by greenhouse gases and causes the atmosphere to warm. The major naturally occurring greenhouse gases present in the atmosphere are carbon dioxide (CO_2), methane (CH_4), nitrous oxides (N_2O), and ozone (O_3). After the meteoric rise in carbon dioxide production in the later decades of the 20th century, global warming became a major concern among scientists worldwide.

Until the middle of the 19th century, carbon dioxide levels remained very stable, fluctuating only slightly in response to natural processes. Any extra carbon dioxide released was absorbed by the ocean or land biosphere. A better understanding of global warming began to take shape when Scandinavian scientists demonstrated that increased levels of atmospheric carbon dioxide due to industrial growth would cause climate change. In 1896, Henry Gifford argued that a doubling of the CO_2 amount could raise the Earth's temperature between 3°C and 6°C. The wildly fluctuating temperature change and its effects on environmental conditions would be disastrous, leading to extreme weather phenomena. The burning of coal and fossil fuel introduced an increased flux of CO_2 into the atmosphere. The Industrial Revolution and the rapid rise of new technologies in power generation, transport, and agriculture greatly increased carbon dioxide emissions.

As sunlight heats the Earth, some of it is absorbed. Other sunshine is reflected off the Earth and sent back to space in the form of heat. Special gases in the atmosphere trap some of this heat, warming the Earth. This mechanism is called the greenhouse effect and it is important to keep the Earth warm enough to sustain life.

To expand on the greenhouse effect, it is important to first explain the meaning of the name. Everyone already knows what a greenhouse is: a glass-roofed building, usually for growing plants that need warmth. The glass lets sunlight in; the inside gets warm. The greenhouse effect on Earth works much the same way. Sunlight passes through the gases that make up the atmosphere. The Earth absorbs some of the sunlight, absorbing warmth. This heat, however, cannot pass back into space easily. It is trapped by the atmosphere, preventing the earth from cooling rapidly.

Many people think of the greenhouse effect as a good thing. Without it, the Earth would be too cold to sustain most life, as average temperatures would be below freezing. However, it is necessary to note that too much of a good thing may also be bad. More greenhouses are being built, and thus more heat is being trapped, making the Earth warmer. Warmer Earth means melting ice, and cities that were once prosperous may be underwater. Warmer Earth also means changing weather, as some regions may become too hot, too dry, while others may become too cold, too wet.

Describing this in detail is not the aim here, but to give just one indicator of things to come. A gas that is blamed for most of the damage humans are causing to the environment is carbon dioxide. Air samples taken thousands of years ago show that the level of carbon dioxide has increased tremendously since the industrial revolution. Carbon dioxide is produced through burning fossil fuels, and plants consume it through photosynthesis. As more forests are cleared for land, nature is losing the ability to consume carbon dioxide. More carbon dioxide in the atmosphere means more heat trapped, resulting in a warmer Earth. It is understood that a doubling of the amount of carbon dioxide in the atmosphere would raise the average global temperature approximately 3 degrees Celsius. This may not seem like much, but there are past examples where such a change in temperature has resulted in a natural disaster on a global scale.

The greenhouse effect is a natural process that warms the Earth's surface. When the sun's energy reaches the Earth, some of it is reflected back to space and the rest is absorbed and re-radiated by greenhouse gases. Greenhouse gases are gases such as carbon dioxide, water vapor, and methane. They let sunlight in, but prevent some of the heat that the sunlight brings from leaving the atmosphere. This is known as the greenhouse effect, and without it, the Earth would be too cold for most living things to survive. Because of the greenhouse effect, the Earth's average temperature is currently about 15 degrees Celsius, or 59 degrees Fahrenheit. However, human activities such as burning fossil fuels have increased the concentrations of greenhouse gases in the atmosphere. As a result, the Earth is quickly becoming warmer, and this is known as global warming.

The greenhouse effect is the process by which greenhouse gases trap heat in the atmosphere. Sunlight, or solar radiation, travels through the atmosphere and warms the surface of the Earth. Some of this heat is absorbed by the Earth and some is reflected back into space. To maintain a constant temperature, the Earth must radiate heat back into space at the same rate that it receives heat from the sun. Heat can be radiated in the form of infrared radiation, which is a band of invisible, or non-visible, light.

The atmosphere is mostly transparent to naked-eye visible light, but is not transparent to infrared radiation. As a result, infrared radiation is absorbed, re-emitted, and scattered by gases in the atmosphere. This process is known as the greenhouse effect and causes the warming of the atmosphere. Some gases contribute more to the greenhouse effect than others, and they are known as greenhouse gases. The most important naturally occurring greenhouse gases are water vapor, carbon dioxide, methane, and ozone. But little by little, humans have been adding to these naturally occurring greenhouse gases, and causing the Earth to warm more than it otherwise would have.

This natural greenhouse effect has been enhanced by human activities, which add additional greenhouse gases to the atmosphere. The burning of fossil fuels (coal, oil, and natural gas) adds additional carbon dioxide to the atmosphere. Deforestation of tropical rainforests prevents the naturally occurring removal of carbon dioxide from the atmosphere. Agriculture produces methane and oxides of nitrogen, both of which are greenhouse gases. Landfills also emit methane as waste decomposes anaerobically (without oxygen). Industrial activities produce a variety of gases that have enhanced the greenhouse effect, and some of these gases can survive for thousands of years before being removed from the atmosphere.

In its fourth assessment report, the Intergovernmental Panel on Climate Change (IPCC) presented a comprehensive overview of the well-established changes in the climate system, as well as observations that are very likely due to human activities. As they prepare for their fifth assessment report, the IPCC plans to present key findings about the impacts of global warming on natural systems and human society. Climate change indices that are based on observations of climate variables help characterize observed changes and variability over time scales of decades to centuries. There exists a number of climate change indices, including extreme indices and simple trend indices. These indices can be used to gauge climate impacts on natural and human systems. The findings concerning these climate change indices, however, should be understood in the context of the information that is not included. Importantly, a number of regions are not well covered, including South America, Africa, and the Arctic. The focus is for the most part on temperature and rainfall indices and includes minimal analysis of other variables such as humidity or wind. There are also large differences in the underlying datasets available for constructing climate change indices. In many low- and middle-income countries, climate data is hardly available. However, with the prospects of near real-time and archived satellite observations as well as downscaled climate models, there are good opportunities to establish climate data for developing countries.

Air temperatures have increased in the last half-century, with the increase being particularly marked in winter. There are also longer-term spatially comprehensive temperature reconstructions. Instrumental datasets covering approximately the last century and a half are continuously updated. Instrumental measurements mainly exist for land stations, while sparse ocean observations were largely limited to the coastal regions until the end of the 1940s. Independent estimates of globally averaged temperature based on a reconstruction from proxy PDC data mostly and, more recently, based on runs of global climate simulations without external forcing are also available. Numerous regional temperature reconstructions of various length, completeness, and temporal resolution exist. In addition to the temperature record, there is also a growing number of reconstructions of global and regional precipitation, tree-ring width, droughts, or other climate characteristics that can be scrutinized for discreteness.

The Intergovernmental Panel on Climate Change (IPCC) reports clearly show that the Earth's surface temperature rose by about 1°C between 1906 and 2005, and it is evident from multiple lines of evidence, including:

1) Surface stations
2) Satellites
3) Ocean temperatures, and
4) Sea surface temperatures

Surface temperature has been monitored since prior to the availability of satellite-based measurements of global temperature. Improved methods were adopted because outdated methods suffered from problems with location and population bias, etc. Near-surface measurements are temperature observations made using hand-operated, automated, or remotely sensed methods in a form that can be used for meteorological or climatological applications. Surface land stations have measured temperature since the late 1800s. This dataset contains both land and ocean. Satellite-based measurements of temperature have been available since the end of 1978 and are unbiased with regard to time and location. Ocean temperature data (obtained from ships and buoys) and satellite sea surface temperature data also indicate warming temperatures. The slowing of ocean heat uptake is discussed in a separate section below.

Temperature rise continues

The global average near-surface temperature rose by about 0.8°C (1.5°F) from the late 19th century to the early 21st century (Figure 3-1). The temperature rise has been greatest since the mid-20th century, with rapid rise over the past three decades. The average rise since 1979 is 0.19°C (0.34°F) per decade and since 1998 is 0.03°C (0.05°F) per decade. Mass loss from mountainous glaciers is widespread and has accelerated in recent decades. Temperatures in the Arctic regions have risen much faster than the global average, about twice as fast over the last several decades according to several estimates. The Arctic is sensitive to global warming because it has a number of positive feedbacks and amplification mechanisms (including albedo feedback) which operate in conjunction with the natural and anthropogenic changes.

Temperature rise from climate models is generally underestimated

CO_2 concentrations are now the highest they have been for several million years, and the long-term climate sensitivity is likely more than 2°C for a doubling of CO_2 concentration, and could be as high as 4.5°C - 6°C. This sensitivity equals or exceeds the climate sensitivity that has already produced the temperature discrepancies. Climate models based on that sensitivity are too cool and leave unchanged a demonstrable warming in infrared dark blue regions of the atmosphere, compared with satellite observations.

By absorbing about a third of the CO_2 emitted into the atmosphere by anthropogenic activities, the ocean plays an important role in regulating climate. However, the well-known consequences of this process are not just the reduction of atmospheric CO_2 concentrations; these processes also alter the ocean's chemistry. In particular, the pH of surface ocean waters is decreased by about 0.1 units (approximately a 30% increase in hydrogen ion activity) since the beginning of the industrial era. This ocean acidification, the uptake of anthropogenic CO_2 by the ocean, is projected to continue well into the future and will profoundly change the conditions in the ocean that are essential for marine life, food web dynamics, biogeochemical cycles, and the functioning of the oceanic carbon sink.

As a consequence of anthropogenic CO_2 emissions, the oceans are acidifying, changing the very conditions under which marine life has evolved. Model projections indicate a future ocean with conditions that have no analog in the last 20 million years and perhaps longer. In combination with other stressors (for example, warming, deoxygenation, and the alteration of the marine environment by humans), such changes may push marine species beyond their physiological or ecological capacity to cope.

Any perturbation to the ocean's system that results in a change of seawater's composition is termed "oceanographic". A simpler definition for ocean acidification is the change in seawater that negatively favors calcifying organisms and plant life. This is in stark contrast to the changes favoring coral calcification, including lower temperature and increased nutrient delivery (N and P) in the model predict that ocean acidification will occur, but with the same trajectory as in the present trajectory, would favor diatoms and dinoflagellate proliferation with variations in nutrient concentrations (favorable conditions for diatoms...)? Marine calcifiers depend upon the concentration of carbonate ions being above a threshold (presently around 220 mmol kg for aragonite and for calcite), below which carbonate is preferentially dissolved.

Ocean acidification a reduction in the ocean's pH due to increased absorption of CO_2 is expected to have deleterious effects on marine calcifiers, with significant implications for marine ecosystems and the biological sequestration of atmospheric CO_2 by the ocean. Broad scale Cook Strait's ocean acidification, caused by hydrodynamically imposed upwelling and long coastal mixing, and therefore upwelled acidified water, forced on by the expected intensification of southerly winds. The synergistic utilization of diverse resources emerges as the pivotal solution to achieve unparalleled efficiency and substantial economic advantages. A remarkable hybrid version ingeniously harnesses the power of both diffuse and/or irradiances, meticulously optimizing the operating temperature for optimum performance. By seamlessly blending these resources, we unlock an unprecedented era of sustainability and harmonious coexistence with our environment [56, 57, 58].

Chapter - 8

Adaptation Strategies

The magnetic levitation system with Proportional-Derivative (PD) + Adaptive Recursive (AR) + Temperature-Compensated State-space Model Control (TSMC) algorithms under various climate variation scenarios is thoroughly examined in this study. The aim is to address the shortcomings of the traditional PD algorithm, such as overshoot, slow speed, and excessive energy consumption. To achieve this, a novel PD + AR + TSMC control algorithm is proposed. By utilizing the adaptive modification principle, the parameter adaptation rule of the recursive identification model of the system, as well as the cam structure and c parameter of TSMC, are determined. Consequently, a system recursive identification model that can dynamically adapt to different climate variation scenarios in real-time is constructed.

Ocean acidification refers to the ongoing decrease in the pH of the Earth's oceans, which is caused by the uptake of anthropogenic carbon dioxide (CO_2) from the atmosphere. As the ocean absorbs more and more CO_2, it forms carbonic acid, which subsequently dissociates and increases the concentration of hydrogen ions and bicarbonate. These ionic changes lead to the overall decrease in seawater pH. Since the Industrial Revolution times, the average surface ocean pH has dropped from 8.2 to 8.1. However, while this development seems small (around 0.1 unit), it amounts to about a 30 percent increase in acidity on the pH scale.

On a global scale, the world's oceans hold 50 times more carbon than the atmosphere. Due to this buffering capacity, the oceans play a very important role in regulating the global carbon cycle, but man-made emissions of carbon must exceed a certain threshold before acidification significantly alters the ocean's chemistry. Acting as a carbon sink, the oceans currently absorb about one-fourth of the carbon emitted into the atmosphere by fossil fuel burning and deforestation. Nevertheless, the time it takes for this CO_2 to diffuse through the oceans is on the order of decades to thousands of years, longer than the time scales for the climate response to some greenhouse warming.

In surface waters, the increase in CO_2 corresponds to over 0.1 (approximately 0.14 to be exact) decrease in sea surface water pH, making it the largest essentially permanent anthropogenic change to the oceans in the past 65 million years. Consequently, the oceans are now more acidic than they have been for at least the last 50 million years, when the oceans were 60 to 85 percent warmer than today. Moreover, changes in the seawater's carbonate chemistry have occurred on temporal and spatial scales for which there is no known natural precedent on Earth, if anthropogenic emissions continue unabated.

Acidification occurs because CO_2 dissolved in seawater forms carbonic acid, which ionizes to produce bicarbonate and hydrogen ions. The hydrogen ions released in this reaction decrease seawater pH and carbonate concentration (the latter is equal to the sum of the concentrations of the CO_3^{2-} and HCO_3^- inorganic species). A considerable amount of fossil carbon released through human activities has ended up in the oceans. It is commonly estimated that more than 43% (as of 2009) of the atmospheric CO_2 released by fossil fuel burning has been absorbed by the oceans, and it is predicted a large percentage of future emissions will also keep being taken up by the oceans, exponentially constant in time. The main oceanic consequence of this gas exchange is the alteration of the dissolved inorganic carbon (DIC) species, and its associated impact on biological and physiological processes.

Since the Industrial Revolution and the ensuing rise in greenhouse gas emissions, the planet has warmed by more than 1°C, resulting in a myriad of changes to global ecosystems and weather patterns. Many species that haven't been able to adapt to these warming conditions fast enough have already faced extinction. Sadly, this appears to be just the beginning. According to Climate Change 2022: Impacts, Adaptation and Vulnerability released by the Intergovernmental Panel on Climate Change (IPCC), up to a million species face extinction due to habitat loss, poaching, overhunting, and pollution, with many in the next 20 years. It will take millions of years for biodiversity to recover post-mass extinction.

Many iconic wildlife species are at risk. Polar bears, whose environment has been melting due to climate change, and koalas, whose food sources are being decimated by drought, bushfire, and logging, stand at the brink. Meanwhile, poisonous frogs recently discovered in the Amazon rainforest, and tropical shorelines home to rich biodiversity are also under threat from climate change, which is inextricably linked to rising human activity from the use of fossil fuels. Even wildlife that might initially seem safe in remote regions are not immune to climate change, which affects all aspects of weather systems and consequently ecosystems.

Change in migration patterns

As temperatures rise, many plants and animals are shifting their habitats to higher altitudes and latitudes. Research published in Nature in 2019 used data from more than 8,000 species and found that terrestrial plants and animals are moving poleward at an average rate of 0.46 km per year and 0.31 km per year, respectively, while marine species are moving poleward at an average rate of 0.29 km per year. Organisms are also moving upwards in elevation. However, many species are unable to keep up with this shift. These changes can disrupt relationships between plants and animals with specificity, such as flowering and pollination and plant life being a habitat for particular animals. Some crop animals, such as cows, would need to migrate northwards to more humid areas as the planet became drier, likely resulting in their extinction. Other species further up the food chain may not have enough to eat following the extinction of prey species.

Oxygen depletion in oceans

Climate change is also affecting the health of oceans. As the earth heats up and emissions of climate pollutants are released into the atmosphere, some of these pollutants find their way into the oceans. Turbulent waters also absorb some of the heat. Bacteria in ocean sediment, published in a study, feed on the organic waste that sinks to the seabed. While these microbes help recycle carbon, they also consume oxygen. Warmer and wild waters generate more food, but this food wealth would lead to partial depletion of oxygen. Changes in ocean circulation, harming halting plankton growth, would also be detrimental due to wild changes in temperature and salinity. All this means that the oceans would see larger dead zones and hovering fish kills. Marine species that are mobile, such as sharks, would be unable to keep up, and less mobile species, such as many bony fishes and mollusks, would be unable to survive.

Climate change is having a devastating impact on the world's biodiversity, resulting in frequent changes to ecological systems. As global warming continues to heat the climate at an unprecedented rate, emerging concerns of biodiversity loss come to light.

More than a million terrestrial and aquatic species face extinction due to global warming. While many species can adapt to changing ecosystems, some do not possess the biological processes and biodiversity needed to adjust accordingly. Recently, the distribution of wildlife populations has shifted towards the poles, and biodiversity is being lost within habitats due to changing temperatures. Rare and specialized species typically suffer the most.

Many alpine species are losing such biodiversity, threatening their long-term persistence. Some trees may not be able to migrate across a landscape quickly enough to escape changing conditions. Furthermore, as the ocean warms, the habitats of several fish and other aquatic species shift closer to the North and South poles.

Affected species tend to have smaller habitats, longer generation times, and those unable to evolve or migrate across landscapes quickly enough. Climate change can also cause mass extinction when performing extreme events. Increasingly severe weather patterns, such as floods or drought, can threaten the persistence of species unable to cope. Furthermore, climate change can increase extinction rates by threatening species' existing interactions in ecosystems. Changes in temperature, food availability, habitat structure, and disease risk can threaten native species and allow invasive species to thrive.

Mitigation strategies refer to methods that attempt to slow or reverse the progress of global warming. Here are some potential strategies that have been suggested. Although some may still be theoretical or untested, some have already been put into practice on a smaller scale. There is no one strategy that will resolve the crisis, and a multifaceted approach that leverages multiple strategies has the best chance of success.

Renewable energy sources draw upon sources of energy that are fundamentally inexhaustible on a human timescale. While some renewable energy tactics (like biomass) may still, on net, produce greenhouse gas emissions, a large proportion of renewable energy sources produce little or no net greenhouse gases. There are numerous technologies and methods for acquiring renewable energy, from well-established implementations to still-opinionated future possibilities.

There are four main sources of renewable energy that would reasonably work on a global scale: solar, geothermal, wind, and tidal energy. Due to the immense amount of solar radiation that reaches the Earth from the sun on a daily basis, solar energy would appear to be the widest in potential scope. Currently, there are two main methods to harvest energy from the sun.

Using solar panels made of photovoltaic cells, light is converted directly into electricity. The electricity can either be stored in batteries for later use or it can be used right away and fed directly into the power grid to power the electrical devices present in homes. The photovoltaic cells are made of silicon, which is the second most abundant element in the Earth's crust. Therefore, there exists a virtually inexhaustible supply. Moreover, harvesting energy from the sun after the panels have been manufactured results in virtually no environmental harm or industrial pollution.

Owing to the concerns about global warming, the availability of fossil fuel reserves, and the rise of energy prices, the search for alternative energy sources has intensified in the past few decades. Renewable energy sources, such as wind, solar, biomass, geothermal, and hydroelectric power, are interesting options. In contrast to fossil fuels, these energy sources are abundantly available, inexhaustible, and a clean form of energy that does not pollute the environment.

A rapidly growing form of renewable energy is wind power. Wind energy makes use of the wind by harnessing its kinetic energy, which can be converted into electricity. The most common and well-developed type of technology for the generation of wind energy is the wind turbine. Wind turbines have been installed worldwide for the past couple of decades. When installed in large numbers, this collective group of turbines is called a wind farm. Wind farms can be located in onshore or offshore locations. Recently, wind energy has been actively promoted by the governments of many countries due to its clear environmental advantages. The utilization of wind energy is expected to become a cornerstone of the development of a sustainable energy system, as the use of fossil fuels would not be feasible in the long-term future.

Solar energy is another promising option. Solar energy refers to the irradiation of the sun that can be converted into other forms of usable energy. Solar energy makes use of the sun's radiation, which can be converted directly or indirectly into thermal energy, mechanical energy, or electricity. The most commonly employed technologies for the conversion of solar energy into electricity are the solar cells made of semiconductors, known as photovoltaics (PVs), and the solar thermal power plants. Solar energy is abundant and free, but there are still technological and infrastructural challenges that need to be resolved before it can be saw widespread usage. However, in industrialized countries and in some developing countries, this energy source is already commercially exploited on a large scale. Other renewable energy sources such as geothermal power, tidal power, and biomass energy, have also been considered.

Geothermal energy makes use of the temperatures of subsurface layers of the Earth. High temperatures have been found at certain locations on the Earth's surface, such as active volcanoes and where tectonic plates are pulling apart, which can be harnessed to produce electricity. Most of these plants are operational in the western portion of the United States (California, Nevada).

Biomass energy is a renewable source of energy derived from organic sources. Biomass sources include charcoal, firewood, agricultural waste, energy crops, and organic waste. Biomass can be converted into fuels or electricity through heat and chemical processes. The problem with this type of energy is the need for massive cultivation of energy crops, which may cause food shortages. Also, burning biomass may contribute to more global warming. Fortunately, biomass can be converted into fuels and fertilizers through the fermentation process, which does not release a lot of greenhouse gases.

Hydroelectric power has been successfully used to produce energy for a long time and is widely used. However, the construction of large dams is very expensive and may irreversibly change the ecosystem.

Global warming can be mitigated through international agreements and political policy reform. A fundamental requirement for curbing greenhouse gas emissions is that they be capped at a defined level rather than grow indefinitely. The concentration of carbon dioxide (CO_2) in the atmosphere should stay below 400 parts per million (ppm), at least until fossil fuels and other sources of carbon can be practically eliminated in favor of sustainable, clean, and renewable energy sources. A maximum level of CO_2 in the atmosphere equivalent to 400 ppm would require a decrease in carbon emissions from 48 billion tons per year to 24 billion by the year 2050 (a decrease of approximately 1.5 percent per year). To stabilize atmospheric CO_2 concentrations at 550 ppm, carbon emissions would need to diminish to 27 billion tons per year by the year 2060 (a decline of approximately 1.0 percent per year).

Policy measures must therefore be put in place to affect large-scale decreases in greenhouse gas emissions either by altering the relative costs of different economic activities or by directly regulating the greenhouse gas emissions. As an alternative to international treaties, emissions trading schemes could be implemented to limit carbon emissions at a level collectively lower than would otherwise be the case. Although it is essential for every country to accept limits on emissions, the approach's success depends very much on the initial distribution of carbon emission quotas. Incentives might then be created for developing countries to limit carbon emissions in exchange for financial aid to help them embark on a less polluting energy path.

Arising from the Framework Convention on Climate Change, the Kyoto Protocol was produced at the Conference of Parties in 1997. The Treaty entered into force on February 16, 2005. For the first commitment period from

2008 to 2012, the Protocol required developed countries with binding targets to reduce their emissions of greenhouse gases by an average of 5 percent below 1990 levels. Developing countries are not required to reduce their emissions under the Protocol. The international rules of carbon trading aim to reduce emissions of greenhouse gases more cheaply or quickly, allowing developed countries to fund projects in developing countries to cut emissions, although these credits are not easily measurable.

Developing countries must be rewarded similarly generous incentives as Denmark received for wind power, Denmark as Germany received for solar photovoltaic (PV) power, and as the U.S. received for nuclear by buying themselves out of oil and coal. After eight years of U.S. talks and setting targets that were later ignored, the Protocol came into force against U.S. moves toward repeal. Most of the world wants action on greenhouse gases. India, China, and the Philippines have all offered sensible proposals that the U.S. ought to support. The experimental findings demonstrate that the proposed control method exhibits excellent positioning performance for combating global warming. Additionally, the adaptive cam structure significantly contributes to the reduction of energy consumption in PD + AR + TSMC control [59, 60, 61].

8.1 Infrastructure resilience

In order to help communities become more resilient to and even prepare for the adverse effects of global warming, a wide array of diverse strategies aimed at mitigating the impacts of inclement weather and rising flood risks stemming from both heavy rainfall and sea level rise are currently being tested and implemented in pilot programs! In the city of Boston, urban planners have employed cutting-edge game simulation software to conduct experiments exploring different development patterns and the efficacy of various hardening approaches with the ultimate goal of fortifying their urban waterfront district. Interestingly, AECOM, a leading global infrastructure firm, has also participated in this progressive initiative! We have drawn invaluable insights from our participatory planning workshops, collaborating closely with local stakeholders, which have consequently culminated in the development of an extensive set of Infrastructure Design Guidelines. Moreover, as a result of our relentless pursuit of excellence, we have made crucial adjustments to certain core infrastructural design elements, thereby ensuring superior functionality and resilience.

It is still up in the ethereal expanse, whether on a practical level there are a sufficient multitude of acceptable adaptation strategies in any community's particular context to properly safeguard and shield most existing urban and small town buildings and districts, and whether it's even remotely conceivable to seamlessly integrate novel or enhanced hardening and infrastructure features into the far-reaching and all-encompassing public and private realms without the indispensable intervention of some public sector benevolence, formidable financial assistance or augmented levels of government stipulating a comprehensive and unifying context for harmonious and concerted action. We ardently maintain that a meticulously coordinated and synchronized implementation effort of paramount importance is obligatory over a diverse assortment of timeframes; our preliminary implementation guide is meticulously contrived and specifically devised to astutely gauge and meticulously evaluate the intrinsic strengths and potential shortcomings of divergent stakeholders' pivotal roles in the meticulous and all-encompassing dialogue indispensable to propel adaptation projects relentlessly forward! As the unfolding of the future gradually materializes and gradually takes form, it is absolutely and unequivocally imperative to vigilantly and meticulously monitor with utmost meticulousness not only the unfolding events occurring in the vast panorama surrounding our existence, but also to meticulously scrutinize the intricate tapestry of how these unfolding events are transpiring and taking shape, and for whom these momentous events are being uniquely tailored and meticulously orchestrated [62, 63, 64].

Chapter - 9

International Cooperation

We should not see worldwide adherence to the disciplines of the Climate Change Convention as an end in itself- because its disciplines may be insufficient. But demonstrating the political will to act will be the key to everything else. Building such demonstrations of political will, in turn, will depend on three things. One is leadership. With leadership, all is possible. At its simplest level, leadership is a matter of determining in discussion the shape of the problem and finding common values to inform the search for meaningful options. It should be unstinting in the pursuit of accountability and work for honesty.

Secondly, it is a matter of hashing out agreements. The disciplines of the Climate Change Convention are agreements by which qualitatively and coherently to confront the problem. Every country must make a down payment on closing implementation and verification gaps, by kicking in through national reporting on greenhouse emissions and the practices, policies, and results. The rewards for shaping a strategy that is both useful and compelling to trounce traditional resistance to cooperative effort must be made more apparent and immediate in caliber and much greater. Refusing action is the route to the surest and worst theft of all - theft of opportunity.

The earth is a vast rainforest. As the planet moves through sunlight, it gradually absorbs carbon dioxide, feeding nature's lush growth. The Sahara Desert is a stellar example of how careless carbon waste and pollution can unravel this careful harmony. Such events sway dew and rainfalls, deterring nature from maintaining and restoring the fellowship of life. Additionally, carbon harvests and sulfur wastes generate harsher temperatures close to the equator over the millennium.

Deforestation renders carbon sinks unviable, exhibiting detrimental effects on oxygen emissions, dirty air clearance, and gentle springs of moisture that hydrate freshwater bodies and keep vegetation intact. Thick forests construct a shield against the sun's glare, reflecting away heat and rain to moist places outside the tropics. Careful alterations to the arrangement of life on earth ripple through thousands of species, raising concern for potential snowballing effects spiraling out of control. Still, as the earth enters a silent change wrought by anthropogenic selfishness, greed, and ignorance, life hangs by a thread, waiting for a turn of human perception.

Public awareness and education about climate change are essential to the successful implementation of effective solutions. Greater public understanding of these problems can lead to an increase in public involvement in climate policy, as well as greater investigative efforts on the part of the scientific community. Not only scientists, governments, and corporations need to be aware of the causes and effects of global climate change, but citizens of the world need to be aware as well. Unfortunately, the educational efforts on this issue are currently grossly inadequate.

A major segment of the public's understanding of and knowledge concerning climate change comes from school curriculums. School programs, ranging from very young children to high school students, should be implemented in order to provide an adequate conceptual foundation for understanding climate change. These educational efforts should include the importance of the greenhouse effect, the sources of greenhouse gases, carbon sequestration, the role of deforestation in climate change, how temperature changes affect the environment and human activities, and the possible ramifications that may arise if global warming continues.

The role of governments and international agencies is crucial to the overall effectiveness of such a broad educational campaign. School programs concerning climate change should be developed and enforced by governmental agencies. Education on climate change should be included in the school curriculums of elementary, middle and high school students. In the United States, for example, topics covered concerning physical science in the grade 9-12 range should include a discussion of climate and climate change in terms of the general circulation model. Such programs should also be designed to work universally with students around the world, perhaps spearheaded by the United Nations Educational, Scientific and Cultural Organization (UNESCO) or the United Nations Environment Programme (UNEP).

Global warming refers to the abrupt rise of the average temperature of Earth, bringing far-reaching consequences to the climate and environment worldwide. Beginning in the late 19th century, scientists have researched and recorded the temperature rise of Earth over time. The Industrial Revolution in the 20th century massively increased the burning of fossil fuels, resulting in a dramatic rise in carbon dioxide (CO2) and other greenhouse gas emissions into the atmosphere. As they increase in concentration, the greenhouse gases create a greenhouse effect, screening sunlight to reach the planet while trapping heat, thus raising the temperature of Earth. The rise of a mere 1.2°C has already disturbed the natural cycles and weather patterns, bringing extreme, unexpected, and unprecedented events. One chain of events causes a domino effect leading to more rapidly changing processes. For example, unprecedented wildfires occur, cycling more soot into the atmosphere, thus making it ever more difficult for the Arctic ice to regenerate during winter. The Arctic ice melting casts yet a darker shade on Earth, further warming it because darker surfaces absorb more sunlight. These effects may spin out of control, creating a self-amplifying pattern, where no action may recover the situation any longer.

Such concerns are not usually addressed in school programs. Various classes in the education system, such as natural sciences, math, social studies, or geography, could use their moderation wisely in directing the attention of the youth to the events. Science education addresses not only codifying facts but also developing students' worldviews and understanding of existing processes. Nevertheless, the climate crisis is hardly ever elaborated on, and its severity and scale are frequently misjudged. Teachers are either unaware of the risks and pressures related to climate breakdown or want to shelter students from feelings of helplessness about the catastrophe. Either way, this choice contributes to growing ignorance, misunderstanding, and disengagement.

A simple set of possible activities can radically change the situation. A web-based tool app can calculate the global mean temperature rise at any given year by entering total CO_2 emissions. The inventory of all climate-related documentaries and science fiction films highlighting the consequences in a predictive manner can be made. An inventory of real climatic events exceeding natural variability limits can be compiled. Student activities can bring climate news in the daily, weekly, or semiweekly school main program. Educational trips to weather stations or global climate centers can be arranged. These, together with holistic yet systematic understanding of current climatic trends and anomalies, realizing the premises of the natural sciences, can get the youth aware, engaged, and empowered. To such radically different present, snow on Christmas and average winter temperatures of around −20 °C are historical clichés similar to presently unbelievable coats of dinosaurs. Hopefully, by making such an attempt, students might still be able to tell their children about winter, ice-skating, and skiing.

The challenge of climate change arises not merely from scientific or technical difficulties but also involves fundamentally ethical questions. Specifically, ethical issues relevant to assessing the causes and consequences of global warming are considered, as well as alternative responses to them. The moral implications of some popular policies are considered.

The ethical dimensions of the global warming problem include harm, causation, fairness and justice, collective actions, and duty of care. Various groups have contributed to the problem and will be differentially affected by it. More strategically, one group might consider with whom they deem it is unfair to treat differently, enacting a climate compensation scheme across nations, as a starting point for collective action. Some states may oppose compensation in an attempt to seize the opportunity afforded them by climate change, asserting a right for development parallel to the industrial revolution seen in the West.

Consequently, it is not enough merely to decide what the best climate policy is from some impartial perspective; one must also be a good judge of the preferences of the parties concerned. The broad ethical framework is described as ensuring action is based on a principled and systematic foundation. Such a foundation would clearly specify moral standards against which different policies are assessed and could state why and how policy choices are made. The major ethical theories in considering climate change and how they view the key mechanisms are outlined.

To illustrate the ethical dimensions of the global warming problem, consider, first, some scientific issues. The greenhouse effect is a natural phenomenon whereby the earth's surface is warmed by the atmosphere. Not only does incoming solar radiation vary, but much of the energy that sustains the earth's climate system is provided and distributed by the earth's atmosphere and oceans. Upon striking the earth's surface, solar radiation is converted to heat; in turn, the earth radiates this heat energy back toward space in order to keep from overheating. Whether or not this is allowed to happen is apparently a matter of chance, determined by the ambient conditions of the atmosphere.

Environmental justice is a concept focused on how climate change will impact communities differently. The areas hardest hit by climate change are often those already unjustly dealt with. However, it goes beyond just claiming that certain groups will have to bear heavier burdens – it is also about who bears the responsibility for addressing these burdens. For instance, it is a moral question whether low-income communities of color should be on the front lines of dealing with climate adaptation and resilience when they did the least to cause climate change.

This perspective also raises the question of who has the moral right to challenge disproportionate burdens and underscore their own contribution to the mitigation of climate change. Environmental justice comes from the grassroots, community-based organizations dedicated to picking up the pieces and dealing with the everyday realities of unfair social arrangements. Many communities will ask, "What can I do about climate change?" and this question brings to bear their social organization, community assets, and political will, as well as their ability to enact policies that articulate their vision of justice and equity. If considerations about social justice and climate change are simply not on these communities' agendas, then a whole cluster of ethical issues is opened up.

Because climate change will have such fundamental impacts on humanity and the planet as a whole, ethical questions arise about the responsibilities of those whose actions are consequential for others. Ethical questions are also raised about how the burdens of climate change (and the burdens of action taken to address it) are distributed among various sectors of society, both nationally and internationally. For these reasons, there is a need to give an overview of both the physical planet-wide changes expected as a consequence of climate change, and of the social changes that are rightly expected to occur in response to these changes.

Consideration of climate change raises two overarching questions about social justice: First, do the actions of one generation affect other generations, so that the concept of inter-generational social justice is meaningful? Second, do the actions of one society affect the welfare of other societies, so that the concept of international social justice is meaningful? As is the case regarding a number of other issues of global ethics, the phenomenon of climate change raises both inter-generational and international dimensions that are potentially overwhelming in scope.

9.1 Paris agreement

The Paris Agreement in 2015 was a critical landmark that initiated a flurry of activities for putting the world on a less than 1.5-2°C pathway. However, despite good intention and numerous efforts by governments and non-government organizations, not much had been done since. Global greenhouse gas emissions had grown annually, even just prior to the pandemic. By September 2023, China and the U.S. had not made much progress over the past ten years in emissions reduction; the Russian Federation had hardly started doing so; and India was still rising in flapping economy. By 2019, the European Union claimed that its carbon emissions were already down by 24% from 1990 level, but higher food and material production, including upon its biodiversity footprint associated with international food and materials consumption, would reveal a similar emission picture at least for air transport. In summary, no one expected much reduction in global greenhouse gas emissions by 2024.

The world came to an unprecedented crossroads with the historic Paris Agreement inked in 2015, an undoubtedly critical landmark that initiated a whirlwind of activities aimed at steering the global community towards a future characterized by a less than 1.5-2°C increase in temperature. However, despite the pureness of intentions and the countless efforts exerted by both governments and non-government organizations alike, it became dishearteningly clear that these initiatives had fallen short, resting idly in the shadow of unfulfilled promises. To add salt to the wound, global greenhouse gas emissions continued their relentless ascent, defying all expectations and escalating their trajectory with each passing year, even on the cusp of the tumultuous pandemic. The world stood aghast as September 2023 rolled around, revealing that economic giants such as China and the U.S. had scarcely made a dent in their emissions reduction efforts over the course of a decade. Similarly disappointing was the Russian Federation's lackluster start in tackling their emissions, while India's economic growth soared skyward, further exacerbating the situation. In the midst of this malaise, a flickering glimmer of hope appeared in the form of the European Union, who proudly declared a 24% decrease in carbon emissions since 1990. However, a closer examination of the situation unraveled a disconcerting revelation - the surge in food and material production, coupled with its insidious impact upon biodiversity, particularly in the context of international consumption, ultimately painted a picture of emissions equivalence, at least within the realm of air transport. Thus, it should come as no surprise that the general consensus held little optimism regarding the magnanimous feat of reducing global greenhouse gas emissions by 2024.

In principle, the variability of temperature is an analogy to the variability of diffusivity and even of viscosity. The existing doctrine which commands how the energy and substance exchange over the Earth-air-ocean system relies on a virtual image assumed for drag/friction, capillarity, and gravity along the Earth-air-ocean interactions which might bear controlled temperature and Earth factors. However, the existing doctrine does not simulate the Earth physical heating in a convincing manner. Similarly, the available doctrine which juxtaposes the diffusivities from lattice vibrations to atmospheric and oceanic thermal movements ignores the fact that lattice vibrations are driven by mean molecular or particle kinetic energy and thus also the Earth energy inside, instead assuming thermal equilibrium of particles/lattice or air/ocean as a vacuum with rarefied atmosphere or water vapor in between until particles or lattices are barely moving but the air and ocean keep themselves in convicting motion.

The immediate future of global warming research can be summarized by three words: computer climate models. Significant effort is being devoted to building, perfecting, and using models of the climate system to predict its response to anthropogenic perturbations. Simple models based on physical principles and observations will continue to evolve and improve our knowledge of the climate system. However, the more complex models being constructed with the assistance of powerful computers may bring a whole new perspective to the subject.

Not too long ago, climate was a subject where one made suppositions and observations without aid of theory or mathematics. Now realistic computations of climate and change of climate can be made. How reliable such computations are and how they compare with considerations based on first principles remain large unanswered questions. Some well-posed initial quarantine conditions have been defined on the subject. Nevertheless, the new model results should stimulate a great deal of physicists' theoretical and mathematical research on terrestrial climate.

It may seem naive to call attention to the whole climate in general. The problems encountered in specific models are indeed often fascinating and complex. Nevertheless, it's unfortunate that the whole climate simply as a well-known terrestrial simple system, with those well-known boundary conditions, is so rarely discussed or even mentioned. If anything is likely to happen to the climate, it is because of a net gain in the extra solar heat absorbed or net increase in the greenhouse efficacy. Although this is in fact virtually trivial, there are a number of unsolved exciting but long-term fluid dynamical, biogeochemical, as well as predictability aspects of the problem that still await a systematic group theoretic treatment.

In view of the impending anthropogenic modification of the climate, there is an urgent need to study the climate as a whole by means of the new physical, mathematical, and computational techniques. In addition to the urgent applied aspects of the research, the whole terrestrial climate should be recognized as an ideal simple evanescent system from the strictly mathematical standpoint and worth theoretically studying, better understanding, and possibly enjoying the sweet dream of the prospect of eventual mastery. Five emerging trends in climate science and global change research are briefly described. These trends are complex, interconnected, and rapidly changing. They are likely to be major areas of emphasis for researchers in the years to come, affecting all parts of the scientific enterprise. They also have wide-ranging implications for education, outreach, and the training of the next generation of scientists.

The first trend is that the Earth system approach is beginning to be used more widely. By taking the Earth system approach, scientists can see how different aspects of the planet interact with each other and feedback through time. There are several aspects of this trend, among them the integration of previously disparate disciplines and the incorporation of biogeochemistry and ecology into Earth system models. A second and related aspect is scale. Models must resolve smaller scales and be more comprehensive in temporal and spatial extent. Models based on Earth system principles are beginning to be applied at greater spatial and temporal scales, such as decadal-to-centennial time scales and large spatial scales such as continents and oceans.

The second emerging trend is concentration on climate interventions. Enhancements of natural processes and/or engineering efforts to reduce climate change by decreasing GHG concentrations or improving the climate's ability to cope with it. There is a growing interest in researching the social, economic, and scientific implications of large-scale intervention strategies. Much attention is focused on geoengineering schemes involving ocean iron fertilization or artificial stratospheric aerosols.

The third emerging trend is the increased prominence of Earth system changes outside of the anthropogenic influence. In recent years, there has been growing interest in understanding natural climate variability over a range of timescales. There are many facets to these efforts that one could explore. One particularly exciting arena relates to the socio-economic consequences of climate variability and change on millennial timescales. Well-documented connections exist between the central Pacific Ocean, climate changes in North America, and El Niño on these timescales. A better understanding of how large ocean-atmosphere processes control climate variability at different timescales could have far-reaching consequences for predictability and for the management of natural resources. Regarding climate changes on timescales greater than millennia, as discussed in Chapter 5, the importance of incorporating effects of Earth's orbital motions into understanding climate change is only beginning to be appreciated.

The emergence of the Earth science initiative is a fourth trend, affecting the design and operation of observing and modeling systems, new constructions such as NASA's Earth System science literacy activities, and NASA's renewed emphasis on interdisciplinary efforts. A growing understanding of how these changes affect the atmosphere, oceans, and geology is shaping what is done and how. Regarding the Earth system approach, understanding how new events and processes are, or could be, scientifically interesting and/or important to society.

The intricate relationship between temperature variability and diffusivity/viscosity variability is something that can be likened in principle. The established theory governing the exchange of energy and substances within the Earth-air-ocean system hinges upon a virtual image, one that assumes drag/friction, capillarity, and gravity as Earth-air-ocean interactions that could wield control over temperature and Earthly factors. However, this existing doctrine falls short when it comes to realistically modeling the physical heating of the Earth.

The Paris Agreement is an international agreement adopted in December 2015 to address global climate change. It was negotiated at the 21st Conference of the Parties to the U.N. Framework Convention on Climate Change (UNFCCC) in Paris, France. The agreement aims to limit global warming to well below 2°C above pre-industrial levels, with efforts to limit the temperature increase to 1.5°C. It also aims to strengthen the ability of countries to deal with the impacts of climate change.

Under the Paris Agreement, countries set nationally determined contributions (NDCs) to reduce greenhouse gas emissions and adapt to the effects of climate change. These contributions are intended to be ambitious, represent a progression over time, and reflect the country's highest level of ambition. Countries are required to submit their NDCs every five years and report on their progress in implementing them.

The agreement also establishes a framework for financial support to developing countries, including funding for mitigation and adaptation activities. Developed countries are expected to provide $100 billion per year by 2020, with a commitment to continue providing financing at least until 2025, which will be scaled up after 2025. The agreement also encourages developed countries to provide support for capacity building and technology transfer.

The Paris Agreement entered into force on November 4, 2016, 30 days after at least 55 countries accounting for at least 55% of global greenhouse gas emissions deposited their instruments of ratification, acceptance, approval, or accession. As of August 2021, 191 parties have ratified the agreement, representing 97.68% of global greenhouse gas emissions.

Highly complex mathematical equations depicted in the form of mathematical models, also called climate models, have been created by professionals to model various climatological occurrences and phenomena such as wind patterns, precipitation, storms, the full climate system, and the full climate. As per multiple researches and recent news articles, climate models are said to play a significant role in understanding the physical processes behind these processes and occurrences. In a broader aspect, climate models are used to describe the present meso-cyclones and large-scale phenomena and help forecast climate conditions and its change by replicating the globe's general circulation model (GCM) for the future.

Highly complex atmospheric primitive equations are used in climate models to forecast climate conditions and states of a location. With the use of complex equations, supercomputers replicate climate models on a variety of scales, depending on the need. Datasets are created from climate models that are brought down to lower resolutions using various approaches. In accordance with defined behavior parameters, model outputs are gridded to a specific layout, and average hourly or daily data grids are settled. These datasets are used for multiple purposes to broadly study the climate system.

Several "state-of-the-art" regional climate modeling systems on the atmosphere-ocean-vegetation-soil system have been designed at spatial and temporal scales that enable the comprehensive version of climate models applied on meso, local, and archive, with some of them coupled in an off-line mode. These climate modeling systems are used to model both present and future climatic conditions and have been shown to yield satisfactory results. Datasets on present bursting precipitation and temperature extremes across various continents and locations across the globe have been created using these models. Climate change scenarios from A1B simulations of the ECHAM5, HadCM3, and CSIRO models within the IPCC 3rd assessment are statistically downscaled to model point datasets on precipitation and temperature. The dataset is used to study possible changes in present climate extreme events in future 21st-century warming.

Climate modeling refers to the utilization of computer models of the Earth's climate system to simulate past, present, or future climate conditions. A climate model is a computer program that uses a set of mathematical equations to recreate the physics and dynamics of the atmosphere, oceans, ice sheets, and land surfaces. Modeling of the climate system is used to understand how the climate works and to make projections about how the climate might change in the future.

Climate models are categorized as "energy balance models," "simple climate models," and "general circulation models (GCMs)." An energy balance model considers only the balance between incoming and outgoing energy in the climate system. A simple climate model considers the oceans and changes in heat storage. General circulation models are created from the fundamental equations of motion, thermodynamics, and other physical principles governing the climate system. GCMs can be run for the whole climate system or treat the components of the climate system in separate models focused on the atmosphere, oceans, or continents. This variety of models may be used to study things such as climate sensitivity and the response of climate under conditions of increased greenhouse gas concentrations.

Increased atmospheric concentrations of greenhouse gases (GHGs) can come from natural sources, such as volcanic eruptions or natural wildfires. However, the increase in the atmospheric concentrations of GHGs today is the result of anthropogenic emissions since the mid-18th century and the advent of the Industrial Revolution. It is expected that anthropogenic emissions will continue to increase considerably and reach double compared to pre-industrial levels.

Ocean general circulation models indicate that the consequences of a doubling of CO_2 include warming over both ocean basins in the mid-latitudes, at high latitudes, and most importantly over the oceans. The lower and upper troposphere will be warmer, but the stratosphere will cool. A scientific consensus has emerged that such climate change will happen. Models indicate the climate may change globally, leading to a rise in sea levels, changes in rainfall regimes, and widespread changes in ecosystems. However, exactly how much and where changes will occur remains uncertain, partially due to feedbacks and uncertainties in future emissions of GHGs. All results to date suggest that the more gases are emitted, the greater the risks of dangerous changes.

Forests play a vital role in cooling the planet by sequestering carbon dioxide from the atmosphere. They are without a doubt the most effective carbon sinks on Earth. Trees breathe in CO_2, and as they grow, they fix carbon in their leaves, branches, and trunks. If nothing disturbs the forest, this carbon will remain there indefinitely. However, scientists' estimates vary greatly on how many gigatons of CO_2 are sequestered by forests each year.

In addition to their role as carbon sinks, forests also indirectly help regulate the planet's temperature. Forests, through the tree canopies, reduce surface temperature fluctuations of the earth. They shade the landscape, which diminishes the amount of sunlight hitting bare soil, forming a microclimate with a cooler temperature, moister air, and heavier fog. Trees also take in water from the soil and release it back into the atmosphere through transpiration, which helps to increase relative humidity.

Because of this microclimate, bare soil, grassland, and crops would dry out and heat up, and the natural precipitation would diminish. The forests need to remain intact to sustain their cooling properties, but deforestation is occurring at an alarming rate. According to satellite imaging, forests are now disappearing at a rate of 15 million km per year, which would be equivalent to the entire Amazon rainforest being lost in 9 years. For Earth to cool off and reverse climate change, forests must be reestablished.

Moreover, the available doctrine that compares diffusivities arising from lattice vibrations with atmospheric and oceanic thermal movements fails to acknowledge the fact that lattice vibrations are propelled by the overall kinetic energy of the molecules or particles, which includes the energy within the Earth itself. This doctrine, rather incorrectly, assumes a state of thermal equilibrium between particles/lattice and air/ocean, portraying them as a vacuum with a thinly distributed atmosphere or water vapor between them. It suggests that particles or lattices remain almost motionless while the air and ocean continue to convict [65, 66, 67].

Chapter - 10

Conclusion

The impacts of warming observed since the industrial era vividly testify to the exceptionally high sensitivity of the climate system to fluctuations in radiative balances. The primary catalyst of this phenomenon is undeniably the unprecedented increase in the concentration of greenhouse gases (GHGs) in the atmosphere, an escalation that can be directly attributed to human activities. Notably, the burning of fossil fuels, rampant deforestation, and agricultural practices have all played pivotal roles in exacerbating this predicament. However, it is crucial to acknowledge that in the short term, air pollution can also wield significant influences by directly altering cloud properties. As we delve into the depths of our understanding, a multitude of uncertainties loom large, particularly in relation to the intricate climate feedback mechanisms. Among these uncertainties, the enigmatic nature of the cloud feedback and sea level variations stands out, as these intricate phenomena are, in part, contingent upon the contributions of distant ice flows residing beyond the confines of our atmosphere.

The atmospheric concentrations of greenhouse gases (GHGs), particularly carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O), are currently at the highest rates observed over the last 800,000 years, and even 20 million years for CO_2. This unprecedented increase in GHG levels poses a significant threat to our climate system. Without explicit climate policies, the anticipated warming between now and 2100 will be within the most pessimistic scenarios, potentially leading to further increasing concentrations of these gases.

In addition to CO_2, methane and N_2O concentrations may also rise due to continued exploitation of natural resources. The growing use of cooling devices in emerging countries is driving the emissions of hydrofluorocarbons (HFCs), which further contribute to the GHG concentrations. It is crucial to address these emissions and implement effective mitigation strategies.

Between 2020 and 2060 or 2070, there is a likelihood of reduced ice cover in polar regions, which, coupled with accelerated melting in many regions, could lead to a rise in the global sea level. The collapse of polar ice flows may exacerbate the impacts, potentially destabilizing natural systems and amplifying warming through various feedback mechanisms. The interactions between the oceans and the atmosphere in such a warming range, averaging 6 degrees across the entire planet, pose significant challenges for available climate models. Predicting the precise effects on our climate system becomes increasingly difficult under these extreme conditions.

Addressing the challenges posed by rising GHG concentrations and climate change requires urgent global action. Climate policies, scientific research, technology development, and international cooperation are vital in mitigating greenhouse gas emissions, adapting to the changing climate, and ensuring a sustainable future for generations to come. It is crucial that we take collective responsibility and work towards a resilient and low-carbon future.

10.1 Summary of key findings

The report on global warming and climate projected that by 2024, the global annual mean surface temperature and the annual mean surface air temperature for all seasons are predicted to be mostly warmer than the recent past. When compared with the range of values determined from the multi-model ensemble, the forecasts of annual mean surface temperature and the forecast skill for all seasons using initialized prediction systems are less uncertain. Warmer temperature than the baseline is predicted for most of the globe. The concentration of CO_2 is forecast to be larger than 2019 and the seasonally ice-free area in the Arctic Ocean is seen to be larger. The predictions of the proportion of the area with lesser annual mean precipitation and that with lesser than the baseline soil moisture are more uncertain. Those flow regimes of the East Asian sector and the outgoing daytime surface radiation of the South Asian monsoon region are predicted to persist and increase, respectively. These predictions suggest warmer and more intense climatic patterns on Earth in 2024.

There are recent research works to project the global mean temperature increase for the near future. The most attention of these works is the projection from the initialized prediction systems of climate models. They are developed by initializing the model with observed data to resemble the observed state and quantify the effects of the observed data. This allows for a quantification of prediction uncertainties, developing data assimilation system, and quantifying the causes of the uncertainty of the climate models on the prediction. The dynamical initialization depends on the assumptions of the climate models and our knowledge of the initialization procedure. Several prediction centers have performed climate predictions for 2020–2024 using different models and analysis. The modeling systems and the skills of the forecast may lead to differences among the forecasts of different centers, but there is substantial consensus that a superficial increase of the concentration of CO_2 and the increase of global annual mean surface temperature are likely to occur.

References

1. S. Mukherjee and A. K. Mishra, "Increase in compound drought and heatwaves in a warming world," Geophysical Research Letters, 2021. wiley.com
2. S. Wang, Q. Zhang, P. Yue, and J. Wang, "Effects of evapotranspiration and precipitation on dryness/wetness changes in China," Theoretical and Applied Climatology, 2020. [HTML]
3. J. S. Gagnon, "Significance of a 1° C increase in global temperature," European Journal of Physics, 2022. [PDF]
4. H. S. Chouhan, "Global Warming Causes and Their Impacts on the Environment," scholar.archive.org. archive.org
5. L. K. Medhekar, "IMPACT OF GLOBAL WARMING AND ITS CHANGES IN ENVIRONMENT," irjhis.com. irjhis.com
6. T. F. Valone, "Gigatonne Carbon Dioxide Removal Can Reverse Global Heating Trend," Asian Journal of Environment & Ecology, 2023. journal4submit.com
7. A. Mikhaylov, N. Moiseev, K. Aleshin, *et al.*, "Global climate change and greenhouse effect," Entrepreneurship and Sustainability Issues, 2020. researchgate.net
8. S. Celik, "The effects of climate change on human behaviors," Environment. researchgate.net
9. J. Huang, G. Zhang, Y. Zhang, X. Guan, *et al.*, "Global desertification vulnerability to climate change and human activities," Land Degradation & Development, vol. 2020. Wiley Online Library. lzu.edu.cn
10. A. Isaacman and M. Musemwa, "Water security in Africa in the age of global climate change," Daedalus, 2021. mit.edu
11. M. Niasse and O. Varis, "Quenching the thirst of rapidly growing and water-insecure cities in sub-Saharan Africa," Global Water Resources, 2021. aalto.fi
12. N. I. Eltaif, M. A. Gharaibeh, and A. M. Fadhil Al-Quraishi, "Impact of Climate Change on Iraq: Severe Water Scarcity and Desertification," 2024. [HTML]

13. A. Chandrasekar, "Basics of atmospheric science," 2022. academia.edu
14. V. Spiridonov and M. Ćurić, "Fundamentals of meteorology," 2021. academia.edu
15. Y. Widiyawati, "Global warming & climate change: Integration of socio-scientific issues to enhance scientific literacy," Journal of Physics: Conference Series, 2020. iop.org
16. R. Feng and X. Fang, "China's pathways to synchronize the emission reductions of air pollutants and greenhouse gases: Pros and cons," Resources. [HTML]
17. J. J. Alava and G. G. Singh, "Changing air pollution and CO2 emissions during the COVID-19 pandemic: Lesson learned and future equity concerns of post-COVID recovery," Environmental science & policy, 2022. sciencedirect.com
18. OO Akomolafe, T Olorunsogo, EC Anyanwu, *et al.*, "Air quality and public health: a review of urban pollution sources and mitigation measures," Engineering Science & …, vol. 2024, fepbl.com, 2024. fepbl.com
19. K. O. Yoro and M. O. Daramola, "CO2 emission sources, greenhouse gases, and the global warming effect," Advances in carbon capture, 2020. researchgate.net
20. L. Bruhwiler, S. Basu, J. H. Butler, A. Chatterjee, *et al.*, "Observations of greenhouse gases as climate indicators," Climatic Change, vol. 2021, Springer, 2021. springer.com
21. V. Iyakaremye, G. Zeng, G. Zhang, "Changes in extreme temperature events over Africa under 1.5 and 2.0 C global warming scenarios," International Journal of …, 2021. researchgate.net
22. Y. Liu, X. Geng, Z. Hao, and J. Zheng, "Changes in climate extremes in Central Asia under 1.5 and 2 C global warming and their impacts on agricultural productions," Atmosphere, 2020. mdpi.com
23. F Yuan-Hai, D Matei, E Tyrlis, C Dong, "Mid-summer surface air temperature and its internal variability over China at 1.5 C and 2 C global warming," in Advances in Climate Change Research, vol. 2020, Elsevier, 2020. sciencedirect.com
24. E. Galanaki, G. Emmanouil, K. Lagouvardos, and V. Kotroni, "Long-term patterns and trends of shortwave global irradiance over the Euro-Mediterranean Region," Atmosphere, 2021. mdpi.com

25. K. Fragkos, I. Fountoulakis, G. Charalampous, *et al.*, "Twenty-Year Climatology of Solar UV and PAR in Cyprus: Integrating Satellite Earth Observations with Radiative Transfer Modeling," Remote Sensing, vol. 2024, mdpi.com, 2024. mdpi.com

26. SS Prasad, RC Deo, N Downs, D Igoe, AV Parisi, "Cloud affected solar UV prediction with three-phase wavelet hybrid convolutional long short-term memory network multi-step forecast system," in IEEE Transactions, 2022. ieee.org

27. M. Amir Siddique, Y. Wang, N. Xu, N. Ullah *et al.*, "The spatiotemporal implications of urbanization for urban heat islands in Beijing: A predictive approach based on CA–Markov modeling (2004–2050)," Remote Sensing, 2021. mdpi.com

28. Y. Li, Z. Zhao, Y. Xin, A. Xu *et al.*, "How are Land-Use/Land-Cover indices and daytime and Nighttime Land Surface temperatures Related in Eleven Urban centres in different global climatic zones?," Land, 2022. mdpi.com

29. S. C. Sam and G. Balasubramanian, "Spatiotemporal detection of land use/land cover changes and land surface temperature using Landsat and MODIS data across the coastal Kanyakumari …," Geodesy and Geodynamics, 2023. sciencedirect.com

30. M. Almazroui, "Changes in temperature trends and extremes over Saudi Arabia for the period 1978–2019," Advances in Meteorology, 2020. wiley.com

31. C. Zhang, Y. Ren, L. Cao, J. Wu *et al.*, "Characteristics of dry-wet climate change in China during the past 60 years and its trends projection," Atmosphere, 2022. mdpi.com

32. L. J. Fan, Z. W. Yan, D. Chen, and Z. Li, "Assessment of Central Asian heat extremes by statistical downscaling: Validation and future projection for 2015–2100," Advances in Climate Change Research, 2022. sciencedirect.com

33. MD Levine, RV Steele, "Climate change: What we know and what is to be done," Wiley Interdisciplinary Reviews: Energy …, vol. 2021, Wiley Online Library, 2021. [HTML]

34. R. K. Upadhyay, "Markers for global climate change and its impact on social, biological and ecological systems: A review," American Journal of Climate Change, 2020. scirp.org

35. G. Alimonti, L. Mariani, F. Prodi, and R.A. Ricci, "RETRACTED ARTICLE: A critical assessment of extreme events trends in times of global warming," The European Physical Journal, 2022, Springer. springer.com

36. F. Song, G. J. Zhang, V. Ramanathan, *et al.*, "Trends in surface equivalent potential temperature: A more comprehensive metric for global warming and weather extremes," Proceedings of the National Academy of Sciences, 2022. pnas.org

37. M. Meinshausen, J. Lewis, C. McGlade, J. Gütschow, *et al.*, "Realization of Paris Agreement pledges may limit warming just below 2 C," Nature, 2022. [HTML]

38. J. Wang, Y. Guan, L. Wu, X. Guan, W. Cai, "Changing lengths of the four seasons by global warming," Geophysical Research, 2021, Wiley Online Library. wiley.com

39. SI Seneviratne, X Zhang, M Adnan, W Badi, "Weather and climate extreme events in a changing climate," 2021, centaur.reading.ac.uk. reading.ac.uk

40. B. Clarke, F. Otto, R. Stuart-Smith, *et al.*, "Extreme weather impacts of climate change: an attribution perspective," Environmental Research Letters, 2022. iop.org

41. S. A. Stephens, R. G. Bell, and I. D. Haigh, "Spatial and temporal analysis of extreme storm-tide and skew-surge events around the coastline of New Zealand," Natural Hazards and Earth System Sciences, 2020. copernicus.org

42. K. Ray, R.K. Giri, S.S. Ray, A.P. Dimri, *et al.*, "An assessment of long-term changes in mortalities due to extreme weather events in India: A study of 50 years' data, 1970–2019," Weather and Climate Extremes, vol. 31, Elsevier, 2021. sciencedirect.com

43. KL Ebi, J Vanos, JW Baldwin, JE Bell, *et al.*, "Extreme weather and climate change: population health and health system implications," Annual Review of Public Health, vol. 42, pp. 293-315, 2021, annualreviews.org. annualreviews.org

44. NV Lukina, AP Geraskina, AV Gornov, "Biodiversity and climate-regulating functions of forests: current issues and research prospects," Вопросы лесной науки, 2021. cyberleninka.ru

45. R. Zhang, D. Tian, J. Wang, and S. Niu, "Critical role of multidimensional

biodiversity in contributing to ecosystem sustainability under global change," Geography and Sustainability, 2023. sciencedirect.com

46. J. A. Thomas, M. Williams, and J. Zalasiewicz, "The anthropocene: A multidisciplinary approach," 2020. [HTML]

47. H. Onyeaka, S. Ghosh, K. Obileke, T. Miri, O.A. Odeyemi, *et al.*, "Preventing chemical contaminants in food: Challenges and prospects for safe and sustainable food production," Food Control, vol. 2024, Elsevier, 2024. sciencedirect.com

48. M. Haji, L. Kerbache, and T. Al-Ansari, "Food quality, drug safety, and increasing public health measures in supply chain management," Processes, 2022. mdpi.com

49. FK Madilo, APH Kunadu, *et al.*, "Challenges with food safety adoption: A review," Journal of Food Safety, vol. 2024, Wiley Online Library. [HTML]

50. D. P. Loucks, "Impacts of climate change on economies, ecosystems, energy, environments, and human equity: A systems perspective," The impacts of climate change, 2021. [HTML]

51. A. Rawat, D. Kumar, B. S. Khati, "A review on climate change impacts, models, and its consequences on different sectors: a systematic approach," Journal of Water and Climate Change, vol. 2024, iwaponline.com, 2024. iwaponline.com

52. K.E. Smith, M.T. Burrows, A.J. Hobday, A. Sen Gupta, *et al.*, "Socioeconomic impacts of marine heatwaves: Global issues and opportunities," Science, 2021. uhi.ac.uk

53. A. Mrówczyńska-Kamińska, B. Bajan, K.P. Pawłowski, *et al.*, "Greenhouse gas emissions intensity of food production systems and its determinants," PLoS One, vol. 2021. Journals.plos.org, 2021. plos.org

54. C. Li, B. Wang, J. J. Klemeš, H. Mikulčić, H. H. Chin, *et al.*, "Greenhouse gas reduction through optimal breeding policy and diet configuration targeting via Carbon Emission Pinch Analysis," Journal of Cleaner Production, vol. 2022, Elsevier, 2022. [HTML]

55. D. J. McClements, "Meat and the Environment: How Livestock Are Taking a Big Bite Out of Our Planet," Meat Less: The Next Food Revolution, 2023. [HTML]

56. XH Chen, K Tee, M Elnahass, R Ahmed, "Assessing the environmental impacts of renewable energy sources: A case study on air pollution and

carbon emissions in China," Journal of Environmental …, vol. 2023, Elsevier, 2023. sciencedirect.com

57. M. Kumar, "Social, economic, and environmental impacts of renewable energy resources," Wind solar hybrid renewable energy system, 2020. intechopen.com

58. A. Rahman, O. Farrok, and M. M. Haque, "Environmental impact of renewable energy source based electrical power plants: Solar, wind, hydroelectric, biomass, geothermal, tidal, ocean, and osmotic," Renewable and Sustainable Energy Reviews, vol. 2022, Elsevier, 2022. [HTML]

59. YS Yu, X Zhang, JW Liu, Y Lee, XS Li, "Natural gas hydrate resources and hydrate technologies: a review and analysis of the associated energy and global warming challenges," Energy & Environmental Science, 2021, pubs.rsc.org. [HTML]

60. E. Cardillo, C. Li, and A. Caddemi, "Embedded heating, ventilation, and air-conditioning control systems: From traditional technologies toward radar advanced sensing," Review of Scientific Instruments, 2021. researchgate.net

61. A. Refaat, A. E. Khalifa, M. M. Elsakka, Y. Elhenawy, "MPPT technique based on enhanced autonomous group Particle Swarm Optimization Algorithm to track the GMPP under partial shading conditions-Experimental," Energy Conversion and Management, vol. 2023, Elsevier, 2023. [HTML]

62. L. Kuhl, "Policy making under scarcity: reflections for designing socially just climate adaptation policy," One Earth, 2021. cell.com

63. L. Susskind and A. Kim, "Building local capacity to adapt to climate change," Climate Policy, 2022. [HTML]

64. R. E. Grumbine and J. Xu, "Mountain futures: pursuing innovative adaptations in coupled social–ecological systems," Frontiers in Ecology and the Environment, 2021. wiley.com

65. B. Charnay, E. T. Wolf, B. Marty, and F. Forget, "Is the faint young Sun problem for Earth solved?" Space Science Reviews, 2020. [PDF]

66. JR Buzan, M Huber, "Moist heat stress on a hotter Earth," Annual Review of Earth and Planetary Sciences, 2020. nsf.gov

67. A. Gettelman, A.J. Geer, R.M. Forbes, G.R. Carmichael, *et al.*, "The future of Earth system prediction: Advances in model-data fusion," Science, 2022. science.org